

FIRST ORDER INVARIANCE OF PLURIGENERA IN POSITIVE CHARACTERISTIC ASSUMING W_2 LIFTING

ANDREW EGBERT

ABSTRACT. This is the end part of what was previously Surf-DIOPDP0.7 which is the Invariance of Plurigenerae in positive characteristic and all kodaira dimensions assuming a lift of $(X, \lceil \Delta \rceil)$. Note that Suh has shown there is not always a lift of $K_X + \lceil \Delta \rceil$ to $W_2(k)$ - the point here is to assume such a lift exists (roughly equivalent to requiring that the Picard Scheme is reduced), and then to show that the plurigenera deform along a curve (at least to first order).

1. PRELIMINARIES

The discussion of deformations will mostly follow the book [Ser] where the theory of deformations, which is generally discussed for compact complex manifolds is extended to an arbitrary algebraically closed field. The underlying field k will therefore always be assumed to be algebraically closed.

1.1. Deformations.

1.1.1. Derivations.

Definition 1. [Abh] A derivation of a ring S with values in an S -module V is a homomorphism $D : S \rightarrow V$ of additive groups such that for all x, y in S , $D(xy) = xD(y) + yD(x)$. The set of all derivations

of S with values in V is an S -module $Der(S, V)$. If $D(x) = 0$ for all x in a subring R of S , then we say that D is an R -derivation of S or a derivation of S/R . The submodule of all derivations of S/R with values in V is denoted by $Der_R(S, V)$. An R -derivation of S (or usually just an R -derivation) is an element of $Der_R(S, S)$. A derivation of S is an element of $Der(S, S)$.

Example 2. Let $R[x_1, \dots, x_n]$ be a polynomial ring. Then ∂_{x_i} , the formal partial derivative with respect to x_i gives an R derivation.

Remark 3. Let R be a ring equipped with a derivation D . We extend D to the total ring of quotients Q of R as follows: If $h \in R$, then $D'(h) = D(h)$. If $h = \frac{u}{v}$ for u, v non-zero-divisors of R , then

$$D(u) = D'(u) = D'(vh) = vD'(h) + hD(v) = vD'(h) + \frac{u}{v}D(v).$$

Subtracting $\frac{u}{v}D'(v)$ from both sides and dividing by v gives

$$D'(h) = \frac{D(u)}{v} - \frac{u}{v^2}D(v) = \frac{vD(u) - uD(v)}{v^2}.$$

So D' is defined by the usual “quotient rule” from calculus existing in the total ring of quotients Q of a ring R . Now letting $u = \frac{u_1}{u_2}$ and $v = \frac{v_1}{v_2}$ be elements of Q such that u_i and v_i are non-zero divisors. Then $D'(u \cdot v) = D'\left(\frac{u_1 \cdot v_1}{u_2 \cdot v_2}\right) = uD'(v) + vD'(u)$ (after expanding) so in fact D' defines a derivation. Thus D' is the unique derivation on the total quotient ring which extends D .

Remark 4. Consider the total ring of quotients Q of a finitely generated k -algebra $S = k[X_1, X_2, \dots, X_n] / (a_1, a_2, \dots, a_m)$ with k a field. If f, g , and $f \circ g \in Q/S$ (invertible elements of the total quotient ring), then by the quotient rule (remark 3), we have

$$D(f(g(x)))' = D\left(\frac{f_1(g(x))}{f_2(g(x))}\right) = D[f(g(x))] \cdot D(g)$$

since expanding $f_i(g(x))$ and using the product and quotient rule gives a chain rule in $k[X_1, \dots, X_n]$. If g is a vector of such elements, then

$$D(f(g^{(1)}(x), \dots, g^{(n)}(x))) = \sum_{j=1}^n \partial_{x^{(j)}} f(g^{(j)}) \cdot D(g^{(j)}(x)).$$

1.1.2. *Deformations.*

Definition 5. [Ser, 1.2.1] Let X be an algebraic scheme. A cartesian diagram of morphisms of schemes

$$\eta : \begin{array}{ccc} X & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \pi \\ \text{Spec}(k) & \xrightarrow{s} & S \end{array}$$

where π is flat and surjective and S is connected, is called a **deformation** of X over S . The deformation is called **first order** if $S = \text{Spec}(A)$ and $A = k[t] / (t)^2$ the ring of dual numbers. The deformation is called **infinitesimal** if $A \in \text{Ob}(\mathcal{A})$, where \mathcal{A} is the category of local artinian k -algebras with residue field k . An infinitesimal deformation is called n^{th} order if $A = k[t] / (t^{n+1}) \in \text{ob}(\mathcal{A})$. A deformation of X over S is called **trivial** if it is isomorphic to the product family $X \times S \rightarrow S$. An infinitesimal deformation of X is called **locally trivial** if every point

$x \in X$ has an open neighborhood $U_x \in X$ such that

$$\begin{array}{ccc} U_x & \longrightarrow & \mathcal{X}|_{U_x} \\ \downarrow & & \downarrow \\ \text{Spec}(k) & \longrightarrow & S \end{array}$$

is a trivial deformation of U_x . All deformations of a nonsingular variety are locally trivial [Ser, 1.2.4].

1.2. Kodaira Spencer Map.

Theorem 6. [Ser, 1.2.9] *Let X be a nonsingular variety. Then there is a 1 – 1 correspondence*

$$\kappa : \{ \text{isomorphism classes of first order deformations of } X \} \rightarrow H^1(X, T_X)$$

called the Kodaira-Spencer correspondence, where $T_X = \text{Hom}(\Omega_X^1, \mathcal{O}_X)$, such that $\kappa(\xi) = 0$ iff ξ is the trivial deformation. The correspondence κ is given as follows: Given a first order deformation

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \text{Spec}(k) & \longrightarrow & \text{Spec}(k[\epsilon]) \end{array}$$

choose an affine open cover $\mathcal{U} = \{U_i\}$ of X such that $\mathcal{X}|_{U_i}$ is trivial for all i . For each index i , we therefore have an isomorphism of deformations

$$\theta_i : U_i \times \text{Spec}(k[\epsilon]) \rightarrow \mathcal{X}|_{U_i}$$

and for each $i, j \in I$ an automorphism

$$\theta_{ij} := \theta_i^{-1}\theta_j : U_{ij} \times \text{Spec}(k[\epsilon]) \rightarrow U_{ij} \times \text{Spec}(k[\epsilon]).$$

$\theta = x + td(x)$ for some $k[\epsilon]$ derivation, and the Kodaira Spencer map takes θ to d . The $\{d_{ij}\}$ forms a Čech cocycle called the Kodaira Spencer class of the deformation.

1.2.1. *Local Definition.* Now consider a first order deformation as above:

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \text{Spec}(k) & \longrightarrow & \text{Spec}(k[\epsilon]) \end{array}$$

with an affine open cover $\mathcal{U} = \{U_i\}$ of X such that $\mathcal{X}|_{U_i}$ is trivial for all i . For each index i , we therefore have an isomorphism of deformations

$$\theta_i : U_i \times \text{Spec}(k[\epsilon]) \rightarrow \mathcal{X}|_{U_i}$$

and for each $i, j \in I$ an automorphism

$$\theta_{ij} := \theta_i^{-1}\theta_j : U_{ij} \times \text{Spec}(k[\epsilon]) \rightarrow U_{ij} \times \text{Spec}(k[\epsilon]).$$

For a given $U_i \cap U_j$, consider the element

$$\rho_{ij} = \sum_{\alpha} \frac{\partial \theta_{ij}^{(\alpha)}}{\partial t} \cdot \frac{\partial}{\partial z_i^{(\alpha)}}$$

where $\frac{\partial}{\partial t}$ denotes the formal derivative with respect to t . This defines a $k[\epsilon]$ derivation and, since $\theta_{ij} = \theta_i^{-1}\theta_j$ and on triple intersections

$\theta_{ij}\theta_{jk}\theta_{jk}^{-1} = 1_{U_{ijk} \times \text{Spec}(k[\epsilon])}$, then using the chain rule (remark 4), $\partial_t \theta_{ij} = \partial_t (\theta_{ik} (\theta_{kj} (z, t), t)) = \partial_t \theta_{kj} + \partial_t \theta_{ij}$ thus

$$\rho_{ij} - \rho_{ji} =$$

and

$$\rho_{ij} + \rho_{jk} - \rho_{ik} = 0$$

so that $\{\rho_{ij}\}$ is a Cech Cocycle giving an element of $H^1(X, T_X)$. By [Ser, 1.2.9], this cocycle uniquely defines the kodaira spencer class of the deformation.

1.3. Obstruction Class. (See [Lev, lemma 1.2]) Consider an infinitesimal deformation n^{th} order deformation defined over $D_{n+1} = \text{Spec}(k[t]/(t^{n+1}))$:

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \text{Spec}(k) & \longrightarrow & D_{n+1} \end{array} .$$

If \mathcal{L} is an invertible sheaf on $X = X_0$, then a deformation of \mathcal{L} over \mathcal{X} is an invertible sheaf \mathcal{L}' on \mathcal{X} such that $\mathcal{L}' \otimes \mathcal{O}_{X_0} \approx \mathcal{L}$.

Suppose that a sheaf L on X_0 has an n^{th} order infinitesimal deformation to $X_n := X_0 \otimes D_n$. Consider the following short exact sequences:

$$0 \rightarrow k \xrightarrow{\times(t^n)} D_n \xrightarrow{\text{mod } (t^n)} D_{n-1} \rightarrow 0$$

and

$$0 \rightarrow D_{n-1} \xrightarrow{\times t} D_n \xrightarrow{\text{mod } t} k \rightarrow 0.$$

As L is invertible, thus locally free, thus flat, then tensoring with L gives the following short exact sequences:

$$0 \rightarrow L_0 \rightarrow L_n \rightarrow L_{n-1} \rightarrow 0$$

and

$$0 \rightarrow L_{n-1} \rightarrow L_n \rightarrow L_0 \rightarrow 0.$$

Note the isomorphism $tL_n \approx L_{n-1}$ so these can be rewritten as

$$0 \rightarrow t^n L_n \rightarrow L_n \rightarrow L_{n-1} \rightarrow 0$$

and

$$0 \rightarrow tL_n \rightarrow L_n \rightarrow L_0 \rightarrow 0.$$

There is a map $f : t^n L_n \rightarrow tL_n$ given by the inclusion $k \rightarrow k[t]/(t^n)$ and another map $g : L_{n-1} \rightarrow L_0$ given by $k[t]/(t^n) \xrightarrow{\text{mod } t^n} k$. Thus there is an induced diagram of long exact cohomology sequences (c.f. [Lev, equation 1]):

$$\begin{array}{ccccc} H^0(X_n, L_n) & \longrightarrow & H^0(X_n, L_{n-1}) & \xrightarrow{\partial} & H^1(X_n, t^n L_n) \\ = \downarrow & & \downarrow g & & \downarrow f \\ H^0(X_n, L_n) & \longrightarrow & H^0(X_n, L_0) & \xrightarrow{\partial} & H^1(X_n, tL_n) \end{array}$$

Let $s \in H^0(X_n, L_0)$. Then $f(\partial(s)) = 0$ implies that $\partial(g(s)) = 0$. If we assume that s already lifts to n^{th} order, then $g(s) = \bar{s} \in H^0(X_n, L_0)$, and hence $\partial(\bar{s}) = 0$, so that \bar{s} is in the image of the bottom left map. This implies $f(\partial(s))$ gives the obstruction to lifting \bar{s} to $H^0(X_n, L_n)$.

1.3.1. *Local Definition.* This discussion follows [Lev, 1.2]. Suppose the situation of subsection 1.3 and assume s cuts out a Cartier divisor. Consider the map

$$\partial : H^0(X_n, L_{n-1}) \rightarrow H^1(X_n, t^n L_n).$$

By Cech cohomology (c.f. [Har]), this map is given by sending the global section $\{s_i\} \in H^0(X_n, L_{n+1})$ to the cocycle defined by

$$\{t^{n+1}\zeta_{ij}(z_j)\} = \{s_i(g_{ij}(z_j, t), t) - \xi_{ij}(z_j, t) \cdot s_j(z_j, t) \bmod t^{n+2}\}.$$

Note that if $n = 0$, then in this case the maps f, g are identity, and hence (since it clearly extends to 0^{th} order) $\partial_1(s) = 0$. So

$$s_i(g_{ij}(z_j, t), t) - \xi_{ij}(z_j, t) \cdot s_j(z_j, t) \equiv 0 \bmod (t^2)$$

and hence, writing $s_i(z_i, t) = \sum_{l=0}^n s_i^l(z_i) t^l$ as a polynomial in t , then the constant term of $\{t\zeta_{ij}(z_j)\}$ must be a coboundary. Thus applying the formal derivative operation $t \cdot \frac{\partial}{\partial t}$ to the above cocycle, we can simplify as follows:

$$s_i(g_{ij}(z_j, t), t) - \xi_{ij}(z_j, t) \cdot s_j(z_j, t) \equiv 0 \bmod (t^2)$$

$$(n+1)t^{n+1}\zeta_{ij} \equiv t \sum_{\alpha} \frac{\partial s_i}{\partial z^{(\alpha)}} \cdot \frac{\partial g_{ij}^{(\alpha)}}{\partial t} + t \frac{\partial s_i}{\partial t} - t \frac{\partial \xi_{ij}}{\partial t} \cdot s_j - t \xi_{ij} \cdot \frac{\partial s_j}{\partial t} \bmod t^{n+1}.$$

The term $t \left(\frac{\partial s_i}{\partial t} - \xi_{ij} \cdot \frac{\partial s_j}{\partial t} \right)$ defines a coboundary, so the above line is equivalent to

$$t \left(\underbrace{\sum_{\alpha} \frac{\partial s_i}{\partial z^{(\alpha)}} \cdot \frac{\partial g_{ij}^{(\alpha)}}{\partial t} - \frac{\partial \xi_{ij}}{\partial t} \cdot \frac{1}{\xi_{ij}} \cdot s_i}_{Ob(s)} \right)$$

so that under the isomorphism $L_n \xrightarrow{\times t} tL_{n+1}$ we must show the vanishing of the term $Ob(s)$ in parentheses.

Claim 7. On $U_i \cap U_j$, there is the identity $-\frac{\partial \xi_{ij}}{\partial t} \cdot \frac{1}{\xi_{ij}} = m \sum_{\alpha} \frac{\partial}{\partial z_i^{(\alpha)}} \left(\frac{\partial g_{ij}^{(\alpha)}}{\partial t} \right)$ for $L = mK_X$, Cartier.

Proof. This follows similarly to [Lev, 1.3], but I argue slightly differently. The transition functions ξ_{ij} for mK_X are given by $-m$ times the determinant of the matrix $A = \left(\frac{\partial g_{ij}^{(\alpha)}}{\partial z_j^{(\beta)}} \right)$. Note that as A is invertible on $U_i \cap U_j$, using the Jacobi identity, (mK_X Cartier implies we can use the chain rule c.f. remark 4)

$$\begin{aligned} & -\partial_t \det(A)^{-m} \cdot \frac{1}{\det(A)^{-m}} \\ &= m \cdot \det(A)^{-m-1} \partial_t \det(A) \cdot \frac{1}{\det(A)^{-m}} \\ &= m \cdot \frac{1}{\det(A)} \cdot \det(A) \operatorname{tr}(A^{-1} \partial_t A). \end{aligned}$$

Note that

$$\frac{\partial g_{ij}^{(\alpha)}}{\partial z_j^{(\beta)}} \cdot \frac{\partial z_j^{(\beta)}}{\partial z_i^{(\gamma)}} = \frac{\partial g_{ij}^{(\alpha)}}{\partial z_i^{(\gamma)}} = \frac{\partial z_i^{(\alpha)}}{\partial z_i^{(\gamma)}} = \delta_{\alpha\gamma}$$

and thus the inverse of A is given by $A_{\beta\gamma}^{-1} = \left(\frac{1}{k} \frac{\partial z_j^{(\beta)}}{\partial z_i^{(\gamma)}} \right)$, so the above becomes (with derivatives first evaluated with respect to t before substitutions are made):

$$= m \sum_{\alpha} \frac{\partial}{\partial t} \frac{\partial g_{ij}^{(\alpha)}}{\partial z_i^{(\alpha)}} = m \sum_{\alpha} \frac{\partial}{\partial z_i^{(\alpha)}} \frac{\partial g_{ij}^{(\alpha)}}{\partial t}.$$

□

By claim 7, $Ob(s)$ can be written as

$$\sum_{\alpha} \frac{\partial s_i}{\partial z^{(\alpha)}} \cdot \frac{\partial g_{ij}^{(\alpha)}}{\partial t} + m \sum_{\alpha} \frac{\partial}{\partial z_i^{(\alpha)}} \frac{\partial g_{ij}^{(\alpha)}}{\partial t}.$$

By [Eg, 2.7], if we add a simple normal crossings boundary Δ , written on the open set U_i as $\gamma_i = \prod_{\alpha \in I_i} z_{i,\alpha}^{q_{\alpha}}$, to L such that $m(K_X + \Delta)$, then $Ob(s)$ becomes

$$\begin{aligned} D(s) &= m \sum_{\alpha} \frac{\partial}{\partial z_{i,\alpha}} \cdot \frac{\partial g_{ij,\alpha}}{\partial t} \cdot s_i \\ &\quad - \sum_{\alpha \in I_i} \frac{\partial g_{ij,\alpha}}{\partial t} \cdot \frac{mq_{\alpha}}{z_{i,\alpha}} \cdot s_i \\ &\quad + \sum_{\alpha} \frac{\partial g_{ij,\alpha}}{\partial t} \cdot \frac{\partial s_i}{\partial z_{i,\alpha}} \end{aligned}$$

in $H^1(X_n, \mathcal{O}_{X_n}(m(K_{X_n} + \Delta_n)))$.

2. MAIN THEOREM

The idea of the proof is to use the arguments of [Lev] which rely only on degeneration of the Hodge-to-de Rham spectral sequence and

not vanishing theorems. As this degeneration holds under certain conditions in characteristic p , and vanishing theorems are known to not necessarily hold, then this approach may be easier. Note that without the lifting assumption, there are actually counterexamples to invariance of plurigenera for elliptic surfaces (see [KU]). The following theorem of Deligne and Illusie is the key ingredient to the proof assuming a lifting actually exists:

Theorem 8. ([DI]) *Let X be a proper smooth variety over a perfect field k of characteristic $p \geq \dim X$ lifting to the ring $W_2(k)$ of the second Witt vectors. Then the hodge to de Rham spectral sequence:*

$$E_1^{ij} = H^j(X, \Omega^i) \implies H_{DR}^*(X/k)$$

degenerates in E_1 .

Remark 9. The above theorem also has a relative version (c.f. [Ber]), so if we want to lift to n^{th} order, it is better to use the version where X/S , where S is a smooth k -scheme.

Proposition 10. *Suppose that X is a smooth proper variety over a perfect field of characteristic $p > k = \dim(X)$. Assume that (X, Δ) is a klt, log smooth pair with a smooth map to an affine curve $\pi : X \rightarrow T$ such that $(X_0, \lrcorner \Delta_0 \lrcorner)$ admits a lifting to W_2 , where X_0 is the fiber over $0 \in T$. Suppose $H^0(m(K_{X_0} + \Delta_0))$ has a smooth section, and*

$m(K_X + \Delta)$ is integral. Then the plurigenera

$$h^0(m(K_{X_t} + \Delta_t))$$

is independent of $t \in T$.

Proof. The proof is along the lines of [Lev] which the author previously generalized to the case of klt pairs in [Eg]. Cover X by coordinate neighborhoods U_i with coordinates (z_i) . Let g_{ij} be transition functions for \tilde{X} satisfying $z_i^{(\alpha)} = g_{ij}^{(\alpha)}$ on X . Let X_0 denote the special fiber and X_n be a scheme flat over $k[t]/(t^{n+1})$ such that $X_n \times_{\text{Spec } k[t]/(t^{n+1})} k \approx X_0$. As in the characteristic 0 case (c.f. [Eg]), suppose s , a smooth section of $h^0(m(K_X + \Delta)/k)$, lifts to $H^0(K_{X_n} + \Delta_n)$ let \bar{s} denote the image of the lift on the central fiber. As in [Eg], the obstruction to lifting $\bar{s} \in H^0(K_{X_0} + \Delta_0)$ to an element of $H^0(X_{n+1}, L_{n+1})$ can be written as

$$\begin{aligned} Ob(s) &= m \sum_{\alpha} \frac{\partial}{\partial z_{i,\alpha}} \cdot \frac{\partial g_{ij,\alpha}}{\partial t} \cdot s_i \\ &\quad - \sum_{\alpha \in I_i} \frac{\partial g_{ij,\alpha}}{\partial t} \cdot \frac{mq_{\alpha}}{z_{i,\alpha}} \cdot s_i \\ &\quad + \sum_{\alpha} \frac{\partial g_{ij,\alpha}}{\partial t} \cdot \frac{\partial s_i}{\partial z_{i,\alpha}} \end{aligned}$$

in $H^1(X_n, \mathcal{O}_{X_n}(m(K_{X_n} + \Delta_n)))$. This is defined through cech cohomology, and hence works in positive characteristic as well. Applying the same calculation as in ([Eg, 4.3]), $D(s) = d\mu$ where μ is a certain element of $H^1(Y_n, \Omega_{Y_n/T}^{k-1})$, where Y_n is the m -fold cover $f : Y_n \rightarrow X_n$ branched over $m(K_{X_0} + \lceil \Delta_0 \rceil)$. Also, by [XieIII, 3.6], Y_0 lifts to W_2

under the assumption that $(X_0, \lrcorner \Delta_0 \lrcorner)$ lifts to W_2 . The Hodge to de Rham spectral sequence on Y_0 has E_1 page

$$\begin{array}{ccc}
 H^0(\Omega_{Y_0}^k) & & H^1(\Omega_{Y_0}^k) \\
 \uparrow & & \uparrow \\
 H^0(\Omega_{Y_0}^{k-1}) & & H^1(\Omega_{Y_0}^{k-1}) \\
 & & \vdots \\
 \uparrow & & \uparrow \\
 H^0(\Omega_{Y_0}^0) & & H^1(\Omega_{Y_0}^0) \quad \dots
 \end{array}$$

and degenerates in E_1 , which means the map $d : H^1(\Omega_{Y_0}^{n-1}) \rightarrow H^1(\Omega_{Y_0}^n)$ is the zero map. Thus, at least in the case that $n = 0$, we can extend s to second order as $d\mu = D(s) = 0$. \square

Remark 11. Note that to extend to higher orders, we must have a relative version of the theorem [XieIII, 3.6].

REFERENCES

- [Abh] Abhyankar, Shreeram Shankar. Lectures on algebra. Singapore: World Scientific, 2006.
- [Ber] Bertin, José. Introduction to Hodge theory. American Mathematical Soc., 2002.
- [Bir] Birkar, Caucher. "Existence of flips and minimal models for 3-folds in char p." arXiv preprint arXiv:1311.3098 (2013).

- [CP] Cossart, Vincent, and Olivier Piltant. "Resolution of singularities of threefolds in positive characteristic II." *Journal of Algebra* 321.7 (2009): 1836-1976.
- [CR] Chatzistamatiou, Andre, and Kay Rülling. "Higher direct images of the structure sheaf in positive characteristic." *Algebra & Number Theory* 5.6 (2012): 693-775.
- [DI] Deligne, Pierre, and Luc Illusie. "Relèvements modulop 2 et décomposition du complexe de de Rham." *Inventiones Mathematicae* 89.2 (1987): 247-270.
- [Ek] Ekedahl, Torsten. "Canonical models of surfaces of general type in positive characteristic." *Publications Mathématiques de l'IHÉS* 67.1 (1988): 97-144.
- [Eg] Egbert, Andrew. "A Note on the Deformation Invariance of Plurigenera for KLT Pairs." preprint (2015)
- [Har] Hartshorne, Robin. *Algebraic geometry*. Vol. 52. Springer Science & Business Media, 1977.
- [HMX2010] Hacon, Christopher, James McKernan, and Chenyang Xu. "On the birational automorphisms of varieties of general type." arXiv preprint arXiv:1011.1464 (2010).
- [KU] Katsura, Toshiyuki, and Kenji Ueno. "On elliptic surfaces in characteristicp." *Mathematische Annalen* 272.3 (1985): 291-330.
- [KK94] Kollár, János, and Sándor Kovács. "Birational geometry of log surfaces." preprint (1994).
- [KM] Kollár, János, and Shigefumi Mori. *Birational geometry of algebraic varieties*. Vol. 134. Cambridge University Press, 2008.
- [Ko] Kollár, János. *Singularities of the minimal model program*. Vol. 200. Cambridge University Press, 2013.
- [Lev] Levine, Marc. "Pluri-canonical divisors on Kähler manifolds." *Inventiones mathematicae* 74.2 (1983): 293-303.

- [Lev-2] Levine, Marc. "Pluri-canonical divisors on Kähler manifolds, II." *Duke Math. J* 52.1 (1985): 61-65.
- [LS] Liedtke, Christian, and Matthew Satriano. "On the birational nature of lifting." *Advances in Mathematics* 254 (2014): 118-137.
- [Sat] Satriano, Matthew. "De Rham theory for tame stacks and schemes with linearly reductive singularities." *arXiv preprint arXiv:0911.2056* (2009).
- [Ser] Sernesi, Edoardo. *Deformations of algebraic schemes*. Vol. 334. Springer Science & Business Media, 2007.
- [Suh] Suh, Junecue. "Plurigenera of general type surfaces in mixed characteristic." *Compositio Mathematica* 144.05 (2008): 1214-1226.
- [Tan] Tanaka, Hiromu. "Minimal models and abundance for positive characteristic log surfaces." *Nagoya Mathematical Journal* (2015).
- [Tan2] Tanaka, Hiromu. "The X-method for klt surfaces in positive characteristic." *arXiv preprint arXiv:1202.2497* (2012).
- [Tan3] Tanaka, Hiromu. "The trace map of Frobenius and extending sections for threefolds." *arXiv preprint arXiv:1302.3134* (2013).
- [Wal] Waldron, Joe. "Finite generation of the log canonical ring for 3-folds in char p ." *arXiv preprint arXiv:1503.03831* (2015).
- [Xie] Xie, Qihong. "Strongly liftable schemes and the Kawamata-Viehweg vanishing in positive characteristic." *Math. Res. Lett* 17.3 (2010): 563-572.
- [XieIII] Xie, Qihong, and Jian Wu. "Strongly liftable schemes and the Kawamata-Viehweg vanishing in positive characteristic III." *Journal of Algebra* 395 (2013): 12-23.