

LOG MINIMAL MODELS FOR ARITHMETIC THREEFOLD GERMS

ANDREW EGBERT

ABSTRACT. In this note, I finish the log minimal model program proofs started in [EGSurfDIOP] and [EgSyz]. In the first note, finite generation of the log canonical ring is proven for Kawamata Log Terminal, log smooth pairs of general type on an algebraic space having two dimensional fibers over a DVR. In the [EgSyz], the minimal model program for such pairs is studied. In this note, I study pairs of any Kodaira dimension.

CONTENTS

1. Intersection Theory on X/R	2
2. Log Resolution	3
3. KLT Kawamata Viehweg Vanishing	5
4. Invariance of Plurigenera Special Case	9
5. Invariance of Plurigenera General Case	14
6. Finite Generation (Big Case)	17
7. Finiteness of Models For Surfaces	20
8. Big Abundance / R	22
9. Cone, Rationality, and contraction / R	25
10. Contractions given by Extremal Rays / R	27
11. Negativity Lemma	28
12. Flips	29

13. Running the Minimal Model Program for General Type	30
14. Reductions To General Case	35
15. Existence of Minimal Models Special Case	37
16. Existence of Minimal Models General Case	39
References	41

1. INTERSECTION THEORY ON X/R

Suppose X/R is a proper algebraic space of relative dimension 2 over a discrete valuation ring R with residue field k of characteristic $p > 0$ and perfect fraction field K (See for example the compact Shimura Varieties mentioned in [Suh, 0.1]).

Lemma 1. [KU, 9.3] *Let $\varphi : X \rightarrow \text{Spec}(R)$ be an algebraic two-dimensional space, smooth, proper, separated, and of finite type over $\text{spec}(R)$, where R is a DVR with algebraically closed residue field k , and field of fractions K . Let X_K (resp. X_k) denote the generic geometric (resp. closed) fibre of φ . If D, D' are divisors on X , then*

$$\begin{aligned} K_{X_k}^2 &= K_{X_K}^2 \\ (K_{X_k} \cdot D_k)_{X_k} &= (K_{X_K} \cdot D_K)_{X_K} \\ (D_k \cdot D'_k)_{X_k} &= (D_K \cdot D'_K)_{X_K}. \end{aligned}$$

If X/R is smooth, then Lemma 1 applies to show that for any divisors C, D extending to both fibers, we can define $C \cdot D = (C|_{X_k} \cdot D|_{X_k})_{X_k}$. On the other hand, if X/R is merely normal and proper, but is actually a scheme, the resolution of singularities, Theorem 3, holds. Thus the intersection theory can be defined as in [Tan, Def 3.1]: $f : X' \rightarrow X$ is a

resolution, and $C.D = f^*C.f^*D$ for two divisors C, D extending to both fibers of Y/R . By properness, this intersection extends by linearity to Weil divisors with \mathbb{Q} or \mathbb{R} coefficients. Numerical equivalence and $N^1(X)_{\mathbb{Q},\mathbb{R}}$ are then defined as usual.

Note the two different hypothesis here: The invariance of plurigenera theorem requires a proper, smooth algebraic space of relative dimension 2, while the existence of minimal models requires the intermediate steps of the minimal model program to have some intersection theory, and thus be schemes. Recall that a Cartier divisor C on a variety is usually called nef if $C.D \geq 0$ for all curves C .

Lemma 2. *Let S be an affine Dedekind scheme and $f : X \rightarrow S$ a projective morphism. Let \mathcal{L} be an invertible sheaf on X such that \mathcal{L}_s is nef for every closed point $s \in S$. Then \mathcal{L} is nef.*

Proof. By [Liu, 5.3.24], the statement holds when nef is replaced by ample. Now, c.f. [Laz, 1.4.10], it suffices to choose an ample divisor H which restricts to X_s , so that $D_s + \epsilon H_s$ is ample for all sufficiently small ϵ . Then $D + \epsilon H$ is ample for all sufficiently small ϵ , and so D is nef. \square

2. LOG RESOLUTION

The following statement of resolution will be used. So for example, if X is an algebraic variety over a number field K then, by [MP, III.2], there is a completed arithmetical variety \overline{X} of dimension $\dim(X) + 1$:

$$\overline{X} \rightarrow \overline{\text{Spec } \mathcal{O}_K}.$$

(Note that a complete DVR is excellent).

Theorem 3. [CP2014, 1.1] *Let X be a reduced and separated Noetherian scheme which is quasi-excellent and of dimension at most three. There exists a proper birational morphism $\pi : X' \rightarrow X$ with the following properties:*

- (i) X' is everywhere regular
- (ii) π induces an isomorphism $\pi^{-1}(\text{Reg } X) \approx \text{Reg } X$
- (iii) $\pi^{-1}(\text{Sing } X)$ is a strict normal crossings divisor on X' .

The following Statement of Log Resolution will be used in the Log Smooth case (i.e. the case with (X, Δ) a pair with X smooth and Δ simple normal crossings.)

Theorem 4. [CP2014, 4.3] *Let S be a regular Noetherian irreducible scheme of dimension three which is excellent and $\mathcal{I} \subset \mathcal{O}_S$ be a nonzero ideal sheaf. There exists a finite sequence*

$$S =: S(0) \leftarrow S(1) \leftarrow \cdots \leftarrow S(r)$$

with the following properties:

- (i) for each j , $0 \leq j \leq r - 1$, $S(j + 1)$ is the blowing up along a regular integral subscheme $\mathcal{Y}(j) \subset \mathcal{S}(j)$ with

$$\mathcal{Y}(j) \subseteq \{s_j \in \mathcal{S}(j) : \mathcal{I}\mathcal{O}_{S(j), s_j} \text{ is not locally principal}\}.$$

- (ii) $\mathcal{I}\mathcal{O}_{S(r)}$ is locally principal.

3. KLT KAWAMATA VIEHWEG VANISHING

In this section, I note that a modification of Tanaka's vanishing theorem [Tan2] gives a version of the (log) Kawamata Viehweg Vanishing in which the multiple of the nef divisor is at least computable. There is a similar recent result in [DF], which is more restrictive (although with a much better constant) in that it requires a smooth variety rather than a normal projective klt pair. First, a theorem of Terakawa:

Theorem 5. [Ter] *Let X be a smooth projective surface over an algebraically closed field of characteristic $p > 0$ and let D be a big and nef Cartier divisor on X . Assume that either*

(1) $\kappa(X) \neq 2$ and X is not quasi-elliptic with $\kappa(X) = 1$; or

(2) X is of general type with

$p \geq 3$ and $(D^2) > \text{vol}(X)$ or

$p = 2$ and $(D^2) > \max\{\text{vol}(X), \text{vol}(X) - 3\chi(\mathcal{O}_X) + 2\}$.

Then

$$H^i(X, \mathcal{O}_X(K_X + D)) = 0$$

for all $i > 0$.

Recall the following covering Lemma:

Lemma 6. *Let X be an n -dimensional smooth variety. Let D be a \mathbb{Q} -divisor such that the support of the fractional part $\{D\}$ is simple normal crossing. Moreover, suppose that, for the prime decomposition $\{D\} = \sum_{i \in I} \frac{b^{(i)}}{a^{(i)}} D^{(i)}$, no integers $a^{(i)}$ are divisible by p . Then there*

exists a finite surjective morphism $\gamma : Y \rightarrow X$ from a smooth variety Y with the following properties:

- (1) The field extension $K(Y)/K(X)$ is a Galois extension.
- (2) γ^*D is a \mathbb{Z} -divisor (it seems the degree of the extension is an m clearing the denominators of the $D^{(i)}$.)
- (3) $\mathcal{O}_X(K_X + \lceil D \rceil) \approx (\gamma_*\mathcal{O}_Y(K_Y + \gamma^*D))^G$, where G is the Galois group of $K(Y)/K(X)$.
- (4) If D' is a \mathbb{Q} -divisor such that $\{D'\} = \{D\}$, then γ^*D' is a \mathbb{Z} -divisor, and $\mathcal{O}_X(K_X + \lceil D' \rceil) \approx (\gamma_*\mathcal{O}_Y(K_Y + \gamma^*D'))^G$.

Remark 7. Several aspects of the proof of the above Lemma will be used in the following.

Applying the above to Tanaka's proof seems to give the following:

Theorem 8. ([Tan2, 2.6] *Weak Effective Kawamata-Viehweg Vanishing Theorem*) *Let X be a smooth projective surface over an algebraically closed field of characteristic $p > 2$. Let N be a nef and big \mathbb{R} -cartier and B a nef and big \mathbb{Q} -divisor whose fractional part is simple normal crossing, whose fractional part has no denominators divisible by the characteristic. Then there exists an r , computable in terms of the intersection numbers of the components of B, N , and K_X , such that*

$$H^i(X, \mathcal{O}_X(K_X + \lceil B \rceil + rN + N')) = 0$$

for every $i > 0$, and every nef \mathbb{R} -Cartier \mathbb{R} -divisor N' such that $rN + N'$ is a \mathbb{Z} -divisor.

Proof. (A slightly different proof than the cited Theorem). The fractional part of $B + rN + N'$ is equal to the fraction part of B when $rN + N'$ is a \mathbb{Z} -divisor. Thus we apply Lemma 6 to obtain a degree m cover $\gamma : Y \rightarrow X$ (independent of r and N') where Y is a smooth surface. Then

$$\begin{aligned}
 & H^1(K_X + \lceil B \rceil + rN + N') \\
 &= H^1(K_X + \lceil B + rN + N' \rceil) \\
 &= H^1(\gamma_*(K_Y + \gamma^*(B + rN + N')))^G \\
 &= H^1(Y, K_Y + \gamma^*B + r\gamma^*N + \gamma^*N')^G.
 \end{aligned}$$

the last term vanishing by Terakawa's Theorem and some $r \gg 0$, as by the proof of [KMM, 1-1-1], the cover can be made general type.

Now I claim it is possible to compute an r such that the last term above is zero using the intersections and degrees of components of $B = \sum a_i \Gamma_i$. It will suffice to show that Y is general type and that $(B + rN)^2 > K_Y^2$, so that Theorem 5 can be applied.

As B is big and nef, there exists an ample divisor A and an effective divisor D such that $B \equiv A - \frac{1}{j}D$ for all $j \gg 0$. Following the proof of [KMM, 1-1-1], let $M = kA$ be very ample (we can compute k by [DF]) and such that m clears the denominators of the components of B and such that $mM - \Gamma_i$ is very ample for all i (again we can compute such an m by [DF]: in the case that $\Gamma_i^2 < 0$, then ensure k is large enough so that $kA \cdot \Gamma_i > -\Gamma_i^2$, take $\Gamma'_i = \Gamma_i + (k-1)K_X$ and by [DF, Theorem 1.2] find (computable) m' such that $m'M - \Gamma'_i = m''M - \Gamma_i$ is very ample. Replace m by m'' , and letting $H_i \in |mM - \Gamma_i|$ for all i (thus

$\frac{1}{m}(\Gamma_i + H_i) \sim M$ is very ample), we then have (again c.f. the proof of [KMM, 1-1-1])

$$\begin{aligned}
 & K_Y \\
 &= \tau^* K_X + (m-1) \left(\sum (\tau^* \Gamma_i)_{red} + \sum (\tau^* H_k^{(i)})_{red} \right) \\
 &= \tau^* K_X + \tau^* \left(\sum_i \left(\Gamma_i + H_i - \frac{1}{m}(\Gamma_i + H_i) \right) \right) \\
 &= \tau^* K_X + \tau^* ((m-1)M)
 \end{aligned}$$

so that Y is general type, and $vol(Y) = [((m+1)k+1)A]^2$, so it suffices to pick r large enough that $(B+rN)^2 > [((m+1)k+1)A]^2$ or, taking $j \gg 0$, such that $(B+rN) > [((m+1)k+1)B]^2$. \square

Now I restate Tanaka's vanishing theorem, with a note on the computability in certain circumstances.

Theorem 9. (*Effective KLT Kawamata-Viehweg Vanishing for normal surfaces c.f. [Tan2, 2.11]*) *Let (X, Δ) be a normal projective klt surface over an algebraically closed field of characteristic $p > 2$, where $K_X + \Delta$ is an effective \mathbb{R} -divisor. Let N be a nef and big \mathbb{R} -cartier \mathbb{R} -divisor. Let D be a \mathbb{Q} -Cartier \mathbb{Z} -divisor such that $D - (K_X - \Delta)$ is nef and big. Then there exists an r_0 ¹ such that*

$$H^i(X, \mathcal{O}_X(D + rN + N^l)) = 0$$

¹By Theorem 8, in the log-smooth case, r_0 is computable in terms of the number of components Δ and in terms of the self intersections of these components, and their multiplicities. Otherwise Tanaka's original statement gives just the existence of an r_0 .

for every $i > 0$, every positive real number $r \geq r_0$, and every nef \mathbb{R} -Cartier \mathbb{R} -divisor N' such that $rN + N'$ is a Cartier divisor.

Proof. The computability follows easily from [Tan2, 2.11] except that Weak Kawamata Viehweg Vanishing in the proof is replaced by Theorem 8. Since, in this note I can apply this theorem when (X, Δ) is log smooth (hence X is smooth) and $\lfloor \Delta \rfloor = 0$, here is the proof in that simple case:

$$\begin{aligned}
 & H^1(D + rN + N') \\
 &= H^1(K_X + D - K_X - \lfloor \Delta \rfloor + rN + N') \\
 &= H^1(K_X + D - K_X + \lceil -\Delta \rceil + rN + N') \\
 &= H^1(K_X + \lceil D - K_X - \Delta \rceil + rN + N')
 \end{aligned}$$

since we can move the integral divisor $D - K_X$ into the round up. This last term vanishes by Theorem 8. The vanishing of H^2 follows easily using Serre Duality. \square

4. INVARIANCE OF PLURIGENERA SPECIAL CASE

Lemma 10. [KU, 9.4] *Let $\varphi : X \rightarrow \text{Spec}(R)$ be an algebraic two-dimensional space, proper, separated, and of finite type over $\text{spec}(R)$, where R is a DVR with algebraically closed residue field k , and field of fractions K . Let X_K (resp. X_k) denote the generic geometric (resp. closed) fibre of φ . If X_k contains an exceptional curve of the first kind e , there exists a DVR $\tilde{R} \supset R$, with residue field isomorphic to k , and a proper smooth morphism $\tilde{\varphi} : \tilde{X} \rightarrow \text{Spec}(\tilde{R})$ of algebraic spaces which is separated and of finite type and a proper surjective morphism*

$\pi : X \otimes \tilde{R} \rightarrow \tilde{Y}$ over $\text{Spec}(\tilde{R})$ such that on the closed fibre, π induces the contraction of the exceptional curve e . Moreover, on the generic fibre, π also induces a contraction of an exceptional curve of the first kind.

Proof. (Follows easily from the cited theorem, although I make a small observation about the extension of residue fields). By [A1, Cor 6.2] $\text{Hilb}_{X/\text{Spec}(R)}$ is represented by an algebraic space \mathcal{H} which is locally of finite type over $\text{Spec}(R)$. Let Y be the irreducible component containing the point $\{e\}$ corresponding to the exceptional curve e on the special fiber. Then $e \approx \mathbb{P}_k^1$ and $N_{e/X} \approx \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}$, so Y is regular at $\{e\}$ and of dimension 1. Since e is fixed in the special fiber, the structure morphism $p : Y \rightarrow \text{Spec}(R)$ is surjective. By [Bo, 2.2.14]², we can find an étale cover $\tilde{R} \supset R$ and a morphism $j : \text{Spec}(\tilde{R}) \rightarrow Y$ over $\text{Spec}(R)$ with $j(\tilde{o}) = \{e\}$ (if R is not already complete, then first extend R to a complete DVR using [Gr, 0.6.8.2,3] so that \tilde{R} is again a DVR). As k is assumed algebraically closed, and $\tilde{R} \rightarrow R$ is unramified, then the extension of residue fields is finite and separable at the closed point of R , and hence an isomorphism of residue fields. Let $\hat{p} : \mathcal{E} \rightarrow \text{Spec}(\tilde{R})$ be the pull-back of the universal family over Y . As the closed fibre \mathcal{E}_0 is a projective line, we may choose the morphism j in such a way that the generic fibre \mathcal{E}_1 of \hat{p} is also a projective line. Moreover, \mathcal{E} can be considered as a smooth closed algebraic subspace of codimension 1 in $\hat{X} = X \otimes \tilde{R}$. By Lemma 1, $-1 = \mathcal{E}_0^2 = \mathcal{E}_1^2$. Hence

²If k is perfect, the proof of the cited lemma seems to go through when $f : X \rightarrow S$ is merely smooth at $x \in X$ and $f : X \rightarrow S$ is locally of finite type.

\mathcal{E}_1 is also an exceptional curve of the first kind. Hence by [A2, Cor 6.11] there exists a contraction morphism $\pi : \hat{X} \rightarrow \hat{X}$ over $\text{Spec}(\tilde{R})$ which contracts \mathcal{E} to a section of $\tilde{\varphi} : \tilde{X} \rightarrow \text{Spec}(R)$ where $\tilde{\varphi}$ is proper, smooth, separated and of finite type over $\text{Spec}(\tilde{R})$. \square

Lemma 11. [KK94, 2.3.5] *Let (S, B) be a log-canonical surface over an algebraically closed field of characteristic $p > 0$. If $C \subset S$ is a curve with $C^2 < 0$ and $C \cdot (K_S + B) < 0$, then $C \approx \mathbb{P}^1$ and it can be contracted to a log-canonical point.*

Theorem 12. [Tan, 4.4, 4.6] *Let X be a projective normal surface and let Δ be an effective \mathbb{R} -divisor such that $K_X + \Delta$ is \mathbb{R} -cartier. Let $\Delta = \sum b_i B_i$ be the prime decomposition. Let H be an \mathbb{R} -cartier ample \mathbb{R} -divisor. Then the following assertions hold:*

- (1) $\overline{NE}(X) = \overline{NE}(X)_{K_X + \Delta \geq 0} + \sum \mathbb{R}_{\geq 0} [C_i]$
- (2) $\overline{NE}(X) = \overline{NE}(X)_{K_X + \Delta + H \geq 0} + \sum_{finite} \mathbb{R}_{\geq 0} [C_i]$
- (3) Each C_i in (1) and (2) is rational or $C_i = B_j$ for some B_j with $B_j^2 < 0$.
- (4) Each C_i in (1) and (2) satisfies $0 < -C_i \cdot (K_X + \Delta) \leq 3$.

Theorem 13. [KK94, 2.3.6 (Log MMP For Surfaces)] *Let (S, B) be a log canonical surface (over an algebraically closed field). There exists a sequence of contractions $f : S \rightarrow S_1 \rightarrow \cdots \rightarrow S_n = S'$ such that S' is log canonical (even plt at every point where f^{-1} is not an isomorphism) and satisfies exactly one of the following conditions:*

- $K_{S'} + f_* B$ is nef

- There exists a $g : S' \rightarrow T$ morphism, such that S' is a birationally ruled surface over the curve T .
- (S', f_*B) is a log Del Pezzo surface.

Remark 14. (Facts about terminal pairs) If $\dim X = 2$ and (X, D) is a terminal pair, then X is smooth [Ko2013]. By [KM, 3.43], terminal singularities are preserved after birational contraction of a $K_X + D$ negative extremal ray, as long as the ray is not a component of D .

Proposition 15. *Let (X, Δ) be a terminal pair of relative dimension 2 over a DVR R with algebraically closed residue field k of characteristic $p > 2$ and perfect fraction field K . Assume that $\mathbb{B}_-(K_X + \Delta) \wedge \Delta = \emptyset$ and that $K_X + \Delta$ is \mathbb{Q} -Cartier with $\kappa(K_{X_k} + \Delta_k) = 2$. Then there exists an m_0 such that for $m > m_0$,*

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

The proof of proposition 15 is given in the following claims.

Claim. Assumptions as above, after passing to an extension R' of R , there is a proper, smooth algebraic space X^{min}/R' and an R' morphism $X \otimes R' \rightarrow X^{min}$ such that both $X_k \rightarrow X_k^{min}$ is obtained by successive blow-downs of (-1) curves and $K_{X_k^{min}} + \Delta_{X_k^{min}}$ is nef.

Proof. As k is algebraically closed of characteristic $p > 0$, then by the Cone Theorem 12,

$$\overline{NE}(X_k) = \overline{NE}(X_k)_{K_{X_k} + \Delta_k \geq 0} + \sum \mathbb{R}_{\geq 0} [C_i]$$

with each C_i is rational or $C_i = B_j$ for some B_j a component of Δ with $B_j^2 < 0$. Under the assumption that $\kappa(K_{X_k} + \Delta_k) = 2$, by Theorem 13 then it must be the case that $C_i^2 < 0$. Thus Theorem 11 implies $C_i \approx \mathbb{P}^1$ and C_i can be contracted to a log-canonical point. By Remark 14 (which applies since $\mathbb{B}_-(K_X + \Delta) \wedge \Delta = \emptyset$) the resulting pair is again terminal, hence smooth, and so it must be the case that C_i was an exceptional curve of the first kind. Therefore it is possible to apply Lemma 10.

By Lemma 10, there is a DVR $\tilde{R} \supset R$ such that $\pi : X \otimes \tilde{R} \rightarrow \tilde{X}$ induces the contraction of C_i on X_k and X_K , and further $\tilde{\varphi} : \tilde{X} \rightarrow \text{Spec}(\tilde{R})$ is proper, smooth, separated, and finite type. Note that after a base change, the extension $\tilde{R} \supset R$ induces a finite extension on residue fields, and since k is algebraically closed, it induces identity on residue fields. Now I need to work with \tilde{X} . \tilde{X}_k is projective, so the same process can be repeated. Each extension $\tilde{R} \supset R$ induces a flat base change on the generic fiber and an isomorphism on the special fiber, and blow-ups are preserved by flat base-change, so there is no problem in extending R . As the Picard number of the special fiber drops each step, there are only finitely many steps. \square

Claim 16. We also have $K_{X_K^{min}} + \Delta_{X_K^{min}}$ nef.

Proof. It suffices to apply Lemma 2 to X_k^{min} , and then restrict to X_K . \square

Claim 17. There is an m_0 such that for $m_0|m$, we have

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

Proof. Suppose first that (X_k, Δ_k) is big. By the claims 4 and 16, we achieve X^{min} such that $K_X + \Delta$ is nef on both X_k^{min} and X_K^{min} . By Theorem 9 applied to the special fiber (and the semicontinuity theorem) there exists an $m_0 \gg 0$ such that for $m_0 < m$ and $i > 0$, we have

$$h^i(m(K_{X_K} + \Delta_K)) = h^i(m(K_{X_k} + \Delta_k)) = 0.$$

Thus, applying the invariance of Euler characteristic [Liu, 5.3.22], and birational invariance of the plurigenera, it follows that for $m > m_0$,

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

□

Remark 18. In fact, if $\Delta = 0$, then, at the end of the above proof, Ekedahl's vanishing Theorem [Ek] can be applied and $m_0 = 2$, in general m_0 can be computed by Theorem 9.

5. INVARIANCE OF PLURIGENERA GENERAL CASE

For technical reasons, I give the terminal case and the klt, log smooth case.

Proposition 19. *Let (X, Δ) be a terminal pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 2$ and perfect fraction field K . Assume that $K_X + \Delta$ is \mathbb{Q} -Cartier and $\kappa(K_{X_k} + \Delta_k) = 2$. Assume that all the prime components of Δ dominate R . Then there exists an m_0 such that for $m_0 | m$,*

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

Proof. Since the hypothesis and conclusion are preserved by base change, then after extending R if necessary, we may assume that k is algebraically closed and that R is complete. Recall that the log-canonical ring of $K_{X_k} + \Delta_k$ is finitely generated by [Tan, 7.1] so $N_\sigma(K_{X_k} + \Delta_k)$ is a \mathbb{Q} -divisor. Let

$$\Theta_k = \Delta_k - \Delta_k \wedge N_\sigma(K_{X_k} + \Delta_k)$$

which is therefore also a \mathbb{Q} -divisor.

Presumably, since the prime components of Δ all dominate R , and since N_σ is a \mathbb{Q} -divisor, then we can find a Θ such that $\Theta|_{X_k} = \Theta_k$. Letting $m \gg 0$ to fit the hypothesis of Proposition 15, and sufficiently divisible such that $m(K_{X_k} + \Theta_k)$ is integral, then by definition of N_σ ,

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_k} + \Theta_k)).$$

Furthermore, by Proposition 15,

$$h^0(m(K_{X_k} + \Theta_k)) = h^0(m(K_{X_K} + \Theta_K)).$$

As $\Theta \leq \Delta$, the theorem follows by semicontinuity. \square

Theorem 20. *Let (X, Δ) be a klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is big, \mathbb{Q} -Cartier, and simple normal crossings over R . Then, there exists an m_0 such that for $m \in m_0\mathbb{Z}^+$,*

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

Proof. Since the hypothesis and conclusion are preserved by base change, then after extending R if necessary, we may assume that k is algebraically closed and that R is complete. To begin, I make the reduction of [HMX2010, 1.6]. As (X, Δ) is log smooth, then replacing (X, Δ) by a blow-up of strata, assume (X, Δ) is terminal. Recall that the log-canonical ring of $K_{X_k} + \Delta_k$ is finitely generated (c.f. [Tan, 7.1]) so $N_\sigma(K_{X_k} + \Delta_k)$ is a \mathbb{Q} -divisor. Let

$$\Theta_k = \Delta_k - \Delta_k \wedge N_\sigma(K_{X_k} + \Delta_k)$$

and let $0 \leq \Theta \leq \Delta$ be a \mathbb{Q} -divisor on X/R such that, by log-smoothness $\Theta|_{X_k} = \Theta_k$. Letting $m \gg 0$ to fit the hypothesis of Proposition 15, and sufficiently divisible such that $m(K_{X_k} + \Theta_k)$ is integral, then by definition of N_σ ,

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_k} + \Theta_k)).$$

Furthermore, by Proposition 15,

$$h^0(m(K_{X_k} + \Theta_k)) = h^0(m(K_{X_K} + \Theta_K)).$$

As $\Theta \leq \Delta$, the theorem follows by semicontinuity. \square

Remark 21. Applying Lemma 48 to the proof of Theorem 20, the minimal models of (X, Δ) and (X, Θ) are actually the same, furthermore, all the additional curves in Δ which appear after blowing up strata

must be -1 curves (since contracting them gives a point of the log-smooth surface X_k). Using these two facts, it is possible to compute m_0 as in Theorem 15.

6. FINITE GENERATION (BIG CASE)

Corollary 22. *Situation as above, after possibly extending R , the canonical ring $R(K_X + \Delta)$ is finitely generated over R .*

Proof. By [Tan], there is $m \gg 0$, such that

$$\bigoplus_{m \geq 0} H^0(ml(K_{X_k} + \Delta_k))$$

is generated in degree 1, so that there is a surjection

$$S^l H^0(m(K_{X_k} + \Delta_k)) \twoheadrightarrow H^0(ml(K_{X_k} + \Delta_k)).$$

Consider the following diagram:

$$\begin{array}{ccc} S^l H^0(m(K_X + \Delta)/R) & \xrightarrow{\alpha} & H^0(ml(K_X + \Delta)/R) \\ \downarrow & & \downarrow \\ S^l H^0(m(K_{X_k} + \Delta_k)) & \twoheadrightarrow & H^0(ml(K_{X_k} + \Delta_k)) \end{array}$$

The vertical maps are surjective by the first part of the theorem (and applying Nakayama's lemma and cohomology and base change to pull back the generators of the k -modules). Now applying the version of Nakayama's Lemma given in [GH, Chapter 5.3], surjectivity of the bottom map implies surjectivity of α , and thus $R(K_X + \Delta)$ is finitely generated over R . \square

Remark 23. It is possible to compute an upper bound for the degree of generation of the log-canonical using Theorem 9. This is explored in more detail in [EgDeg].

Definition 24. Let $(X, \sum S_i)$ a log smooth, log-canonical projective pair where X is a two dimensional, normal variety over a perfect field of characteristic $p > 0$. with S_i distinct prime divisors, $V = \sum \mathbb{R}S_i \subset \text{Div}_{\mathbb{R}}(X)$. Define the following sets, the first of which is clearly a polytope.

$$\mathcal{L}(V) = \left\{ \sum_{i=1}^n a_i S_i \in V \mid a_i \in [0, 1] \right\}$$

$$\mathcal{E}(V) = \{ \Delta \in \mathcal{L}(V) \mid K_X + \Delta \sim_{\mathbb{R}} D \geq 0 \}.$$

Given $f : X \rightarrow Y$ a birational contraction, let $\mathcal{C}_f(V)$ denote the closure of $\mathcal{L}(V)$ of

$$\{ \Delta \in \mathcal{E}(V) \mid f \text{ is a log terminal model of } (X, \Delta) \}.$$

C.f. [CL2013, 2.13], (the statement is for characteristic 0 in any dimension by [SC, 3.4], but in the surface case here merely relies on [Tan, 0.2]) then there are birational contractions $f_i : X \rightarrow Y_i$ such that $\mathcal{C}_{f_i}(V), \dots, \mathcal{C}_{f_k}(V)$ are rational polytopes and

$$\mathcal{E}(V) = \bigcup_{i=1}^k \mathcal{C}_{f_i}(V)$$

so that $\mathcal{E}(V)$ is also a polytope.

Proposition 25. [H] *Let X be a two dimensional \mathbb{Q} -factorial normal variety over a perfect field of characteristic $p > 0$. Let $\{(X, \Delta_i)\}_{i \in \{1, \dots, k\}}$*

be a big, log smooth, \mathbb{Q} -Cartier KLT pair for each i and such that $K_X + \sum \Delta_i$ has simple normal crossings support. Then the ring

$$R(X, K_X + \Delta_1, \dots, K_X + \Delta_k)$$

is finitely generated.

Proof. I sketch a proof based on [CL2013, 2.18] which applies in the KLT case (Hashizume's Theorem seems to give the more general, log-canonical case). Let $\sum \text{supp}(\Delta_i) = \sum S_i$ for S_i distinct prime divisors, $D_i = \ell(K_X + \Delta_i + A)$ and $\mathcal{M} = \sum_{i=1}^p \mathbb{Z}D_i$. Let $M^{(N)} = \sum_{i=1}^p \mathbb{Z}ND_i$. As in the cited theorem, there is a constant M' and generators of $\mathcal{M} \cap \mathbb{R}_+(K_X + \mathcal{C}_i)$ for all i of the form $M'(K_X + B)$ for some $B \in \mathcal{L}(V)$. Applying abundance ([Tan]), there is an $M \in \mathbb{N}$ such that for all i, j where B_{ij} span each \mathcal{C}_i , such that $M(f_i)_*(K_X + B_{ij})$ is basepoint free. Thus, c.f. [CZ, 2.11], we are reduced to proving the following: Let (X, B_i) a klt pair for each i and $L_i = a_i(K_X + B_i)$ has basepoint free linear system for each i , (we can assume here that $K_X + B_i$ is big). Let $G = \sum_{i=1}^k b_i L_i$ for positive integers b_i , and assume that $b_\ell > n + 1$. Then the natural map

$$H^0(X, \mathcal{O}_X(G)) \otimes H^0(X, \mathcal{O}_X(L_\ell)) \rightarrow H^0(X, \mathcal{O}_X(G + L_\ell))$$

is surjective. The rest follows the proof of [CZ, 2.7], but substituting the Vanishing Theorem 9 when necessary for the usual Kawamata Viehweg Vanishing. \square

Corollary 26. *Let $\{(X, \Delta_i)\}_{i \in \{1, \dots, n\}}$ be a collection of klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta_i$ is big, \mathbb{Q} -Cartier, and either simple normal crossings over R for each i , or terminal and with components dominating $\text{Spec } R$, and such that $K_X + \sum \Delta_i$ has simple normal crossings support. Then the ring*

$$R(X, K_X + \Delta_1, \dots, K_X + \Delta_n)$$

is finitely generated over R .

Proof. C.f. the proof of [CZ, 2.11], it suffices to prove that if $G = \sum_{i=1}^k m_i (a(K_X + \Delta_i))$, with fixed $a \in \mathbb{N}$, $m_1, \dots, m_k \geq 0$, and $l \in \{1, \dots, l\}$ then there exists m_l such that the following maps are surjective

$$\begin{aligned} H^0(X, \mathcal{O}_X(G - a(K_X + \Delta_i))) \otimes H^0(X, \mathcal{O}_X(a(K_X + \Delta_i))) \\ \rightarrow H^0(X, \mathcal{O}_X(G)). \end{aligned}$$

By applying Nakayama's Lemma and Theorem 20 and Proposition 19 as in the proof of Theorem 22, it suffices to show the surjectivity on the central fiber which holds by proposition 25. \square

7. FINITENESS OF MODELS FOR SURFACES

The following theorem is not anything new, but is stated here as the minimal model program is not usually run with scaling on a surface (in [KK94, Tan], termination is proven without scaling).

Theorem 27. *Let X be a two dimensional normal variety over a perfect field of characteristic $p > 0$. Let (X, Δ) be a log smooth KLT pair and A an ample \mathbb{Q} -divisor on X with a good minimal model (which exists for non-negative Kodaira dimensions by [Tan]). Then there exists an $\epsilon > 0$ such that the minimal models (and the output of the minimal model program with scaling) for $(X, \Delta + tA)$ are all isomorphic for $t \in [0, \epsilon]$.*

Proof. (Essentially same as [Lai, 26,27] and simplifies due to the fact that a birational isomorphism in codimension 1 on a surface gives an isomorphism). Let $\phi : (X, \Delta) \rightarrow (X_g, \Delta_g)$ be a good minimal model. Let $f : X_g \rightarrow Z = Proj R(K_{X_g} + \Delta_g)$ the contraction (which exists by since $K_{X_g} + \Delta_g$ is nef, hence semiample by [Tan]). Then ϕ contracts the divisorial part of $\mathbb{B}(K_X + \Delta)$. Pick $t_0 > 0$ such that $(X_g, \Delta_g + t_0\phi_*A) = (X_g, \Delta_g + t_0A_g)$ is klt with A ample on X (note A_g is big, not in general nef). For H ample on X_g , let $\psi : X_g \rightarrow X'$ be the minimal model program with scaling, which terminates (again by [Tan]), and gives a minimal model of $(X_g, \Delta_g + t_0\Delta_g)$ over Z . For any curve contracted by f , $(K_{X_g} + \Delta_g) \cdot C = 0$, hence $K_{X'} + \psi_*\Delta_g = K_{X'} + \Delta' \equiv_Z 0$. Thus curves contracted by ψ have trivial intersection with $K_{X_g} + \Delta_g$, and intersect negatively with A_g . Thus changing t does not affect which curves intersect negatively with $K_{X_g} + \Delta_g + tA_g$, and so X' is a minimal model of $(X_g, \Delta_g + tA_g)$ for all $t \in (0, t_0]$.

Now $\Delta' + t_0A'$ with $A' = \psi_*A_g$ (which is big) implies by the Theorem 12, that there exist only finitely many $K_{X'} + \Delta' + t_0A'$ negative

extremal rays in $\overline{NE}(X')$. These are all necessarily just intersecting A' negatively, so decreasing t_0 , eventually we get to a point where a further decrease in t_0 doesn't change the number of negative extremal rays. Pick such a t_0 (Now we are on X' , not over Z). Again shrinking t_0 , suppose that $\psi \circ \phi$ is discrepancy negative w'r't $(X, \Delta + tA)$ for all t in the half open interval $(0, t_0]$. Note that $\mathbb{B}(K_X + \Delta + t_0A) \subset \mathbb{B}(K_X + \Delta)$, so ψ contracts only the curves not contracted by ϕ . Thus $\psi \circ \phi$ is discrepancy negative on the closed interval $[0, t_0]$, so X' is a minimal model of $(X, \Delta + tA)$ for all $t \in [0, t_0]$. Thus $\mathbb{B}(K_X + \Delta + tA)$ has the same divisorial components for all $t \in [0, t_0]$. Hence, any two minimal models for different $t \in [0, t_0]$ are birational and isomorphic in codimension one, and as the total dimension is two, they are thus isomorphic. \square

8. BIG ABUNDANCE / R

Definition 28. Let D_1, \dots, D_r \mathbb{Q} -Cartier \mathbb{Q} -divisors on X . D_i are called **adjoint divisors** on X if they are of the form $D_i = K_X + \Delta_i$ for some **pair** (X, Δ_i) where X is normal and projective, $\Delta_i \geq 0$ and is a \mathbb{Q} -divisor. Let

$$R = R(X; D_1, \dots, D_r) = \bigoplus_{(n_1, \dots, n_r) \in \mathbb{N}^r} H^0(X, n_1 D_1 + \dots + n_r D_r).$$

and define the **support** of R to be

$$\text{Supp } R = \left(\sum_{i=1}^r \mathbb{R}_+ D_i \right) \cap \text{Div}_{\mathbb{R}}^{\text{eff}}(X).$$

Theorem 29. [CL, 3.5] *Let X be a normal scheme of relative dimension 2, proper over S , the spectrum of a DVR which has perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Let D_1, \dots, D_r be \mathbb{Q} -Cartier divisors on X . Assume that $R(X; D_1, \dots, D_r)$ is finitely generated. Let $D : \mathbb{R}^r \ni (\lambda_1, \dots, \lambda_r) \mapsto \sum \lambda_i D_i \in \text{Div}_{\mathbb{R}}(X)$ be the tautological map.*

(1) *The support of R is a rational polyhedral cone.*

(2) *If $\text{Supp } R$ contains a big divisor and $D \in \sum \mathbb{R}_+ D_i$ is pseudoeffective, then $D \in \text{Supp } R$.*

(3) *There is a finite rational polyhedral subdivision $\text{Supp } R = \bigcup C_i$ such that σ_{Γ} is a linear function on C_i for every geometric valuation Γ of X . Furthermore, there is a coarsest subdivision with this property, in the sense that, if i and j are distinct, there is at least one geometric valuation Γ of X such that (the linear extensions of) $\sigma_{\Gamma}|_{C_i}$ and $\sigma_{\Gamma}|_{C_j}$ are different.*

(4) *There is a finite index subgroup $\mathbb{L} \subset \mathbb{Z}^r$ such that for all $n \in \mathbb{N}^r \cap \mathbb{L}$, if $D(n) \in \text{Supp } R$, then $\sigma_{\Gamma}(D(n)) = \text{mult}_{\Gamma}(D(n))$.*

Proof. (1, 2) follow easily from the cited theorem. For 3, 4 note that [ELM, 4.7] is stated for an arbitrary Noetherian scheme, and this theorem is the only result used in the proof of this result which is given in [CL, 3.5]. □

Theorem 30. ([CL, 3.6]) *Let (X, Δ) be a klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is \mathbb{Q} -Cartier that*

$R(K_X + \Delta, K_X + \Delta + A)$ is finitely generated. If $K_X + \Delta$ is pseudo-effective, then it is \mathbb{Q} -effective.

Proof. As in the source paper. □

Let Γ a geometric valuation on X . Define

$$\sigma_\Gamma = \inf \{ \text{mult}_\Gamma D' \mid D \sim_{\mathbb{R}} D' \geq 0 \} \in \mathbb{R}.$$

As a result of Theorem 26, there is the following abundance theorem:

Corollary 31. [CL, 3.8] *Let (X, Δ) be a klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $\mathcal{R} = R(K_X + \Delta, K_X + \Delta + A)$ is finitely generated for an ample \mathbb{Q} -divisor A on X . Assume that Δ is big. If $K_X + \Delta$ is nef, then it is semi-ample.*

Proof. $\text{Supp } \mathcal{R}$ contains a big divisor $(K_X + \Delta + A)$. Thus, any pseudo-effective

$$D \in \mathbb{R}_+(K_X + \Delta) + \mathbb{R}_+(K_X + \Delta + A)$$

is in $\text{Supp } \mathcal{R}$. By Theorem 29(2),

$$\text{Supp } \mathcal{R} = \mathbb{R}_+(K_X + \Delta, K_X + \Delta + A).$$

For any $\epsilon > 0$, and any geometric valuation Γ on X , as $K_X + \Delta + \epsilon A$ is ample,

$$\sigma_\Gamma(K_X + \Delta + \epsilon A) = 0$$

so that σ_Γ is zero on $\text{Supp } \mathcal{R}$, and thus the centre of Γ is not in $\mathbb{B}(K_X + \Delta)$ for any such Γ . □

Corollary 32. *Let X be a normal projective variety of dimension 2 (over a perfect field k) and let D_1, \dots, D_r be \mathbb{Q} -Cartier \mathbb{Q} -divisors on X . Assume that the ring $R = R(X; D_1, \dots, D_r)$ is finitely generated, and let $\text{Supp } R = \bigcup C_i$ be a finite rational polyhedral subdivision such that for every geometric valuation Γ of X , σ_Γ is linear on C_i , as in Theorem 29. Denote $\varphi : \sum \mathbb{R}_+ D_i \rightarrow N^1(X)_\mathbb{R}$ the projection, and assume there exists k such that $C_k \cap \varphi^{-1}(\text{Amp } D) \neq \emptyset$. Then $C_k \subset \text{Supp } R \cap \varphi^{-1}(\text{Nef } X)$. If the subdivision is coarsest, then $C_k = \text{Supp } R \cap \varphi^{-1}(\text{Nef } X)$*

Proof. As in the source, but applying Theorems 29 and 31. □

9. CONE, RATIONALITY, AND CONTRACTION / \mathbb{R}

Definition 33. [CL, def 4.1] Let W a finite dimensional real vector space, $C \subset W$ a closed convex cone spanning W , and $v \in W$. The visible boundary of C from v is

$$V = \{w \in \partial C \mid [v, w] \cap C = \{w\}\}.$$

Corollary 34. (c.f. [CL, 4.2]) (*Kawamata's Rationality, Cone and Contraction Theorem*) Let (X, Δ) be a pair of relative dimension 2, proper over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $v_0 = K_X + \Delta$ is \mathbb{Q} -Cartier and is obtained from the log smooth klt pair (X', Δ') after a finite number of steps of the minimal model program with scaling associated to a big divisor A with $K_X + \Delta + tA$ nef on X/R for some $t > 0$. Let V be the visible boundary of $\text{Nef}(X/R)$ from $v_0 \in N^1(X)_\mathbb{R}$ of the divisor $K_X + \Delta$ and $u_0 \in V \cap ([K_X + \Delta], [K_X + \Delta + tA])$. Then there exists a

locally polyhedral neighborhood U of u_0 with $U \subset V$ such that $\forall u' \in U$, u' is semiample.

Proof. This is essentially a local version of the cited theorem. Let $\mathcal{C} = \mathbb{R}_+ v_0 + \text{Nef}(X)$. Choose ϵ such $B(u_0, \epsilon)$ is a sup-norm ball about u_0 with rational vertices $w_j = K_X + \Delta + A_j$, $j = 1, \dots, r$ which are klt and pull back to big log smooth klt pairs on X' , and such that ϵ is small enough that any $K_X + \Delta + A' \in B(u_0, \epsilon)$ pulls back to a log smooth divisor on X' . Let \mathcal{B} denote the convex hull of $B(u_0, \epsilon)$. Then by Theorem 26, the ring

$$R = R(X, K_X + \Delta + A_1, \dots, K_X + \Delta + A_r)$$

is finitely generated. Let $\varphi : \sum \mathbb{R}_+(K_X + \Delta + A_j) \rightarrow N^1(X)_{\mathbb{R}}$ the natural projection so that $\varphi(\sum \mathbb{R}_+(K_X + \Delta + A_j)) = \mathbb{R}_+ \mathcal{B}$. Theorem 29 implies that $\text{Supp } R$ is a rational polyhedral cone which intersects the interior of the nef cone. Therefore, by Theorem 29(2), if U is the portion of $\partial \text{Nef}(X/R)$ contained in $B(u_0, \epsilon)$, then $\varphi^{-1}(U) \subset \text{Supp}(R)$. Let $\text{Supp } R = \bigcup \mathcal{L}_k$ be the coarsest subdivision given by Theorem 29(3). For some k , $\mathcal{L}_k \cap \varphi^{-1}(\text{Amp } X) \neq \emptyset$, so by Theorem 32 $\mathcal{L}_k = \text{Supp } R \cap \varphi^{-1}(\text{Nef}(X))$, so that U is locally polyhedral. If $u' \in U$, with divisor D then there exists an ample \mathbb{Q} -divisor A' such that $D \sim_{\mathbb{Q}} K_X + \Delta + A'$. By choice of ϵ , D pulls back to a log smooth divisor on X' , and so that $R(K_X + \Delta + A', K_X + \Delta + A' + A'')$ is finitely generated for any ample A'' such that $D + A'' \in B(u_0, \epsilon)$. Thus Theorem 31 gives that u' is semiample. \square

Lemma 35. [CL, Lemma 5.1] *Let X and Y be \mathbb{Q} -factorial projective schemes smooth over a DVR R with algebraically closed residue field. Let $f : X \rightarrow Y$ be a birational contraction, and let $\tilde{f} : \mathbb{K}(Y) \approx \mathbb{K}(X)$ the induced isomorphism on function fields. Then:*

- (1) $f_* \operatorname{div}_X \varphi = \operatorname{div}_Y f(\varphi)$ for every $\varphi \in \mathbb{K}(X)$;
- (2) for every geometric valuation Γ on $\mathbb{K}(X)$, and for every $\varphi \in \mathbb{K}(X)$, we have $\operatorname{mult}_\Gamma(\operatorname{div}_X \varphi) = \operatorname{mult}_\Gamma(\operatorname{div}_Y \tilde{f}(\varphi))$;
- (3) if f is an isomorphism in codimension one, then $f_* : \operatorname{Div}_{\mathbb{R}}(X) \rightarrow \operatorname{Div}_{\mathbb{R}}(Y)$ is an isomorphism, and for every $D \in \operatorname{Div}_{\mathbb{R}}(X)$, there is an isomorphism $H^0(X, D) \approx H^0(Y, f_* D)$.

Proof. Follows easily from the cited Theorem. □

Lemma 36. [CL, 5.2] *Let X, Y be proper of relative dimension 2, over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K and $f : X \rightarrow Y$ a birational map which is an isomorphism in codimension one. Let $\mathcal{C} \subset \operatorname{Div}_{\mathbb{R}}^{\operatorname{eff}}(X)$ be a cone, and fix a geometric valuation Γ of X . Then the asymptotic order of vanishing σ_Γ is linear on \mathcal{C} if and only if it is linear on $f_* \mathcal{C} \subset \operatorname{Div}_{\mathbb{R}}^{\operatorname{eff}}(Y)$.*

Proof. As in the cited theorem, substituting Lemma 35 as necessary. □

10. CONTRACTIONS GIVEN BY EXTREMAL RAYS / R

Lemma 37. [CL, 6.2] *Let (X, Δ) be a log smooth klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic*

$p > 0$ and perfect fraction field K . Suppose (X, Δ) is not nef, and A is a big \mathbb{Q} -divisor such that $(X, \Delta + A)$ is klt, and $K_X + \Delta + A$ is nef. Let λ be the nef threshold. Then $\lambda \in \mathbb{Q}^+$, and there is $R \subset \overline{NE}(X)$ with $(K_X + \Delta + \lambda A) \cdot R = 0$ and $(K_X + \Delta) \cdot R < 0$.

Proof. As in the cited theorem, but substituting Theorems 32 and 26 for their corresponding versions. \square

11. NEGATIVITY LEMMA

Lemma 38. [HK, 1.7] *Let $f : X \rightarrow Y$ and $g : X \rightarrow Z$ be birational contractions where all spaces considered are schemes of relative dimension 2 over DVR with perfect residue field. Suppose $f^*(A) + E = g^*(B) + F$ for A ample, B nef, E f -exceptional, and F g -exceptional all divisors extending to both fibers. Then $f \circ g^{-1} : Z \rightarrow Y$ is regular.*

Proof. It suffices to apply negativity lemma (c.f. [Xu, 2.1] for positive characteristic and $\dim \leq 3$) to the special fiber so that $E = F$ and thus $f^*(A) = g^*(B)$. The result then follows from [Ke, 1.0]. \square

Lemma 39. [CL, 6.4] *Let (X, Δ) a projective klt pair, and $f : X \rightarrow Y$ a composite of $(K_X + \Delta)$ -divisorial contractions and $(K_X + \Delta)$ -flips. Then for every resolution*

$$\begin{array}{ccc}
 W & \xrightarrow{p} & X \\
 \downarrow q & \nearrow f & \\
 Y & &
 \end{array}$$

of f ,

$$p^*(K_X + \Delta) = q^*(K_Y + f_*\Delta) + E$$

with $E > 0$ a q -exceptional divisor. Therefore f cannot be an isomorphism. By the above formula,

$$H^0(X, K_X + \Delta) \approx H^0(Y, K_Y + f_*\Delta).$$

Proof. Follows easily from [KM, 3.38]. □

Lemma 40. [Har, Exc II.4.2] *Let X be a reduced scheme, Y a separated scheme, and let f and g be two morphisms from X to Y . Assume that $f|_U = g|_U$ on a Zariski dense open subset $U \subset X$. Then $f = g$.*

12. FLIPS

Definition 41. [KM, 3.33] Let X a normal scheme and D a \mathbb{Q} -divisor on X such that $K_X + D$ is \mathbb{Q} -Cartier. A $(K_X + D)$ -**flipping contraction** is a proper birational morphism $f : X \rightarrow Y$ to a normal scheme Y such that $Exc(f)$ has codimension at least two in X and $-(K_X + D)$ is f -ample. A normal scheme X^+ together with a proper birational morphism $f^+ : X^+ \rightarrow Y$ is called a $(K + D)$ -**flip** of f if

(1) $K_{X^+} + D^+$ is \mathbb{Q} -Cartier, where D^+ is the birational transform of D on X^+

(2) $K_{X^+} + D^+$ is f^+ -ample, and

(3) $Exc(f^+)$ has codimension at least two in X^+ .

The induced rational map $\phi : X \rightarrow X^+$ is sometimes called a $(K + D)$ -flip by abuse of notation.

Theorem 42. *Let (X, Δ) be a projective \mathbb{Q} -factorial klt big pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Suppose that $f : X \rightarrow Y$ is a $K + \Delta$ flipping contraction. If $R(K_X + \Delta)$ is finitely generated, then the $(K + \Delta)$ -flip of f exists.*

Proof. Follows easily as in [KM, 6.2], [CL, 6.3]. □

Remark 43. By [JKW, 1.2], the flip of f in the situation above is \mathbb{Q} -factorial.

13. RUNNING THE MINIMAL MODEL PROGRAM FOR GENERAL TYPE

First a useful lemma due to Gallego and Purnaprajna:

Lemma 44. ([GP], 1.4.1) *Let E and L_1, \dots, L_r be coherent sheaves on a variety X . Consider the map*

$$H^0(E) \otimes H^0(L_1 \otimes \cdots \otimes L_r) \xrightarrow{\psi} H^0(E \otimes L_1 \otimes \cdots \otimes L_r)$$

and the maps

$$H^0(E) \otimes H^0(L_1) \xrightarrow{\alpha_1} H^0(E \otimes L_1),$$

$$H^0(E \otimes L_1) \otimes H^0(L_2) \xrightarrow{\alpha_2} H^0(E \otimes L_1 \otimes L_2)$$

...

$$H^0(E \otimes L_1 \otimes \cdots \otimes L_{r-1}) \otimes H^0(L_r) \xrightarrow{\alpha_r} H^0(E \otimes L_1 \otimes \cdots \otimes L_r).$$

If $\alpha_1, \dots, \alpha_r$ are surjective, then ψ is also surjective.

Theorem 45. *Let (X, Δ) be a projective \mathbb{Q} -factorial klt pair of relative dimension 2 over the spectrum of a DVR $S = \text{Spec } R$ where R has perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Suppose Δ is big and log smooth over R . Then the minimal model program for (X, Δ) can be run, resulting in a (possibly non-terminating) sequence of flips and divisorial contractions.*

Proof. If necessary, extend R to be complete with algebraically closed residue field k . I begin as in [CL, 6.2]. Let A be a big divisor such that $K_X + \Delta + A$ is nef. Let α_1 be the smallest positive real number such that $K_X + \Delta + \alpha_1 A$ is nef. Denote by $\varphi : \text{Div}_{\mathbb{R}}(X) \rightarrow N^1(X)_{\mathbb{R}}$ the natural projection, and let $\|\cdot\|$ any norm on $N^1(X)_{\mathbb{R}}$. Pick finitely many big \mathbb{Q} -divisors H^1, \dots, H^r such that:

- (1) $\|\varphi(\Delta + \alpha_1 A) - \varphi(H^i)\| \ll 1$ for all i
- (2) writing $\mathcal{C} = \sum_{i=1}^r \mathbb{R}_+(K_X + H^i) \subset \text{Div}_{\mathbb{R}}(X)$, we have $K_X + \Delta + \alpha_1 A \in \text{int } \mathcal{C}$, and the dimension of the cone $\varphi(\mathcal{C}) \subset N^1(X)_{\mathbb{R}}$ is $\dim N^1(X)_{\mathbb{R}}$
- (3) (X, H_i) is klt and log smooth for all i with $\text{supp}(K_X + \sum H_i)$ simple normal crossings.

In the first step, since H_i are log smooth, then by Theorem 26, the ring

$$R(X, K_X + \Delta, K_X + H^1, \dots, K_X + H^r)$$

is finitely generated. Thus applying Theorem 32, $\text{Supp } R = \bigcup C_j$ is rational polyhedral and $C_j = \text{Supp } R \cap \varphi^{-1}(\text{Nef}(X))$ is rational polyhedral, and there is a rational codimension 1 face of $\text{Nef}(X)$. Thus $\alpha_1 \in \mathbb{Q}_+$, and we can find an extremal ray $\mathfrak{R} \subset \overline{NE}(X)$ dual to this

face which will satisfy $(K_X + \Delta + \alpha_1 A) \cdot \mathfrak{R} = 0$ and $(K_X + \Delta) \cdot \mathfrak{R} < 0$. By Theorem 31, (after applying Theorem 26 to satisfy the hypothesis) this ray can be contracted under a birational contraction: $f' : X \rightarrow X'$.

If f' is a divisorial contraction, write $f_1 = f'$ and set $\Delta_2 = f_{1*}\Delta$, $A_2 = f_{1*}A$, so A_2 is again big and $K_{X_2} + \Delta_2 + \alpha_1 A_2$ is nef. Note that $R(X_2, K_{X_2} + \Delta_2, K_{X_2} + \Delta_2 + A_2)$ is again finitely generated and $R(X_2, K_{X_2} + \Delta_2)$ is finitely generated since for this is a contraction, and for instance. Pick a new set $\{H_2^i\}$ of big, log smooth H_i satisfying the above hypothesis, (since the pushforwards may no longer be log smooth). Then properties (1)-(3) hold. Thus we can repeat the process using finite generation of

$$\begin{aligned} & R(X_2, K_{X_2} + \Delta_2, K_X + H_2^1, \dots, K_X + H_2^r), \\ & \approx R(X, K_X + \Delta, K_X + H^1, \dots, K_X + H^r). \end{aligned}$$

If, on the other hand, f' is a small contraction, then by Theorem 42, and applying Theorem 22, the flip of the contraction exists (c.f. definition 41). Write f_1 for the induced morphism to the flip and set $\Delta_2 = f_{1*}\Delta$, $A_2 = f_{1*}A$, so A_2 is again big and $K_{X_2} + \Delta_2 + \alpha_1 A_2$ is nef. By Lemma 35, we have finite generation of

$$\begin{aligned} & R(X_2, K_{X_2} + \Delta_2), \\ & R(X_2, K_{X_2} + \Delta_2, K_{X_2} + \Delta_2 + A_2) \end{aligned}$$

and, in small contraction case, by Lemma 35 we can let $H_2^r = f_{1*}H^r$, and finite generation of

$$R(X_2, K_{X_2} + \Delta_2, K_{X_2} + H_2^1, \dots, K_{X_2} + H_2^r)$$

so we can similarly repeat the process. \square

The next theorem assumes we have arrived at a sequence of flips.

Theorem 46. *Notation as above, there is no infinite sequence of flips starting at a given step j of the minimal model program described in Theorem 45. Thus in this case, the minimal model program with scaling ends in a minimal model $(X_g, \Delta_g) = (X_{j'}, \Delta_{j'})$ on which all of the following rings are finitely generated: $R(K_{X_{j'}} + \Delta_{j'})$,*

$$R(K_{X_{j'}} + \Delta_{j'}, K_{X_{j'}} + \Delta_{j'} + A_{j'}),$$

and

$$R_{j'} = R(K_{X_{j'}} + \Delta_{j'}, K_{X_{j'}} + H_{j'}^1, \dots, K_{X_{j'}} + H_{j'}^r).$$

Proof. If necessary, extend R to be complete with algebraically closed residue field k . Suppose to the contrary there is an infinite sequence of flips starting at step j . As in Theorem 45, we have for the j^{th} step, finitely many big \mathbb{Q} -divisors H_1^j, \dots, H_r^j such that:

$$(1) \|\varphi(\Delta_j + \alpha_j A_j) - \varphi(H_j^i)\| \ll 1 \text{ for all } i$$

$$(2) \text{ writing } \mathcal{C}^j = \mathbb{R}_+(K_{X_j} + \Delta_j) + \sum_{i=1}^r \mathbb{R}_+(K_{X_j} + H_j^i) \subset \text{Div}_{\mathbb{R}}(X_j),$$

we have $K_{X_j} + \Delta_j + \alpha_j A_j \in \text{int } \mathcal{C}_j$, and the dimension of the cone $\varphi(\mathcal{C}^j) \subset N^1(X_j)_{\mathbb{R}}$ is $\dim N^1(X_j)_{\mathbb{R}}$

$$(3) (X_j, H_j^i) \text{ are Kawamata Log Terminal}$$

(4)

$$R(K_{X_j} + \Delta_j),$$

$$R(K_{X_j} + \Delta_j, K_{X_j} + \Delta_j + A_j),$$

and

$$R_j = R(K_{X_j} + \Delta_j, K_{X_j} + H_j^1, \dots, K_{X_j} + H_j^r)$$

are all finitely generated.

(5) $K_{X_j} + \Delta_j + \alpha_j A_j$ is nef.

I proceed almost verbatim as in [CL, 6.5]. By construction, C^j contains an open neighborhood of the nef divisor $K_{X_j} + \Delta_j + \alpha_j A_j$, so C^j contains ample divisors in its interior. Thus the cone $\varphi(\text{Supp } R_j) \subset N^1(X)$ also has dimension $\dim N^1(X)_{\mathbb{R}}$. Let $\text{Supp } R = \bigcup_k C_k^j$ the coarsest finite rational polyhedral subdivision from Theorem 29(3). For $i > j$, let $C_k^i \subset \text{Div}_{\mathbb{R}}(X_i)$ denote the proper transform of C_k^j and $C^i \subset \text{Div}_{\mathbb{R}}(X_i)$, the proper transform of C^j . By Lemma 36, for every geometric valuation Γ the asymptotic order function σ_{Γ} is linear on each C_k^j .

By construction if $0 < \alpha \leq \alpha_1$, then $K_{X_j} + \Delta_j + \alpha A_j \in \text{Int } C^j$, so $K_{X_i} + \Delta_i + \alpha_i A_i \in \text{int } C^i$ for all $i > j$. Since $K_{X_i} + \Delta_i + \alpha_i A_i$ is nef, then by requirement (4) above and Lemma 30, $K_{X_i} + \Delta_i + \alpha_i A_i \in \text{Supp } R_i$. Thus by Theorem 32, for each i there exists an index k such that the image of C_k^i in $N^1(X_i)_{\mathbb{R}}$ is a subset of $\text{Nef}(X_i)$. Therefore,

$$\varphi(C_k^1) \subset (f_{i-1} \circ \dots \circ f_1)^* \text{Nef}(X_i).$$

Since there are finitely many cones C_k^j , there are two indices p and q such that $(f_{p-1} \circ \cdots \circ f_1)^* \text{Nef}(X_p)$ and $(f_{q-1} \circ \cdots \circ f_1)^* \text{Nef}(X_q)$ share a common interior point. Thus, by Lemma 38, the map $X_p \rightarrow X_q$ is a morphism. By Lemma 40, it is therefore an isomorphism. This contradicts Remark 43. \square

14. REDUCTIONS TO GENERAL CASE

Lemma 47. [HMX2014, 2.8.3] *Let (X, Δ) be a log smooth pair which is a scheme of dimension at most 3, with the coefficients of Δ belonging to $(0, 1]$, and with X projective. If (X, Δ) has a weak log canonical model, then there is a sequence $\pi : Y \rightarrow X$ of smooth blow ups of the strata of Δ such that if we write $K_Y + \Gamma = \pi^*(K_X + \Delta) + E$, where $\Gamma \geq 0$ and $E \geq 0$ have no common components, $\pi_*\Gamma = \Delta$ and $\pi_*E = 0$ and if we write*

$$\Gamma' = \Gamma - \Gamma \wedge N_\sigma(Y, K_Y + \Gamma),$$

then $\mathbb{B}_-(Y, K_Y + \Gamma')$ contains no strata of Γ' . If Δ is a \mathbb{Q} -divisor, then Γ' is a \mathbb{Q} -divisor.

Proof. Follows easily from the cited theorem. \square

Lemma 48. [HMX2014, 5.3] *Let X/R a projective scheme of relative dimension 2 over a DVR R , with algebraically closed residue field and perfect fraction field. Let (X, Δ) be a log smooth dlt pair with X \mathbb{Q} -factorial and projective and Δ a \mathbb{Q} -divisor. If Φ is a \mathbb{Q} -divisor such that*

$$0 \leq \Delta - \Phi \leq N_\sigma(X, K_X + \Delta),$$

then steps of the $K_X + \Phi$ -minimal model program are steps of the $K_X + \Delta$ minimal model program, and so termination for (X, Φ) implies termination for (X, Δ) .

Proof. (Similar proof to the cited Theorem). Let $f : X \rightarrow Y$ the result of running a $(K_X + \Phi)$ minimal model program and let $Y \rightarrow Z$ be the ample model of $K_X + \Phi$. Let $\Delta_t = t\Delta + (1 - t)\Phi$ so that if $0 < t \ll 0$, then f is also a $(K_X + \Delta_t)$ -MMP. If C generates a $K_X + \Delta_t$ -negative extremal ray, suppose that $(K_X + \Phi) \cdot C > 0$. Applying Theorem 12 on the special fiber,

$$-4 \leq (K_X + \Delta) \cdot C < 0.$$

Then, following the inequalities in [HMX2014, 5.1], gives a contradiction if $0 < t < \frac{1}{5}$. Thus every step of the $K_X + \Delta_t$ with scaling of an ample divisor is $(K_X + \Phi)$ -trivial, so after finitely many steps there is a model $g : X \rightarrow W$ contracting the components of $N_\sigma(X, K_X + \Delta_t)$ (this holds with no changes from [HMX2014, 2.7.1]). Thus, as $\text{supp}(\Delta - \Phi) \subset \text{supp} N_\sigma(X, K_X + \Delta) = \text{supp} N_\sigma(X, K_X + \Delta_t)$, then $g_*(K_Y + \Phi)$ is semiample whenever $g_*(K_X + \Delta)$ is, and [HMX2014, 2.7.2] (which holds using only the log smooth resolution) implies that g is a minimal model of (X, Δ) , which is good when (X, Φ) has a good minimal model. \square

Lemma 49. [GL, 2.3] *Let X/R a scheme which is projective of relative dimension 2 over either a DVR R with algebraically closed residue field and perfect fraction field or over a field. Let (X, Δ) be a log smooth klt pair and $\varphi : W \rightarrow X$ a log resolution of (X, Δ) . Choose Δ_W so that*

$K_W + \Delta_W = \varphi^*(K_X + \Delta) + E$ with Δ_W and E effective \mathbb{Q} -Weil divisors with no common component. Let $F = \sum_{F_i \text{ } \varphi\text{-exceptional prime divisor}} F_i$ and $\Delta_W^\epsilon = \Delta_W + \epsilon F$, then (W, Δ_W^ϵ) is an ϵ -log smooth model of (X, Δ) . Then for $0 < \epsilon \ll 1$, a (good) minimal model $\phi : (W, \Delta_W^\epsilon) \rightarrow (W_{min}, \Delta_{W_{min}}^\epsilon)$ is also a (good) minimal model for (X, Δ) .

Proof. Follows easily from [BCHM, 3.6.10, 3.6.11].

□

15. EXISTENCE OF MINIMAL MODELS SPECIAL CASE

Lemma 50. [HMX2010, 3.1] *Let (X, Δ) be a \mathbb{Q} -factorial klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$ contains no non-canonical centres of (X_k, Δ_k) . Let $f : X \rightarrow Y$ be a step of the $(K_X + \Delta)$ -MMP. If f is birational and V is a non-canonical centre of (X, Δ) , then V is not contained in the indeterminacy locus of f , V_0 is not contained in the indeterminacy locus of f_0 , and the induced maps $\phi : V \rightarrow W$ and $\phi_k : V_k \rightarrow W_k$ are birational, where $W = f(V)$. Let $\Gamma = f_*\Delta$. Then $\mathbb{B}_-(Y_0, K_{Y_0} + \Gamma_0)$ contains no non-canonical centres of (Y_0, Γ_0) (so we may repeat the process), and if V is a non-klt centre or $V = X$, then $\phi : V \rightarrow W$ and $\phi_0 : V_0 \rightarrow W_0$ are birational contractions. If f is a Mori fibre space, then f_0 is not birational.*

Proof. (Follows easily from the cited Theorem). Suppose f is birational. Let V be a non-canonical centre of (X, Δ) . Let $g : X \rightarrow Z$ be the contraction of the extremal ray associated to f (so that $f = g$

unless f is a flip). Let $Q = g(V)$, and let $\psi : V \rightarrow Q$, be the induced morphism. As every component of V_k is a non-canonical centre of (X_k, Δ_k) , then by hypothesis, components of V_k are not contained in $\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$, thus ψ_k is defined at each such component, hence is birational. By upper-semicontinuity of the fibers of ψ , ψ is birational, and thus $\phi : V \rightarrow Q \rightarrow W$, and ϕ_k are both birational.

Now suppose V is a non-klt centre or $V = X$, the above holds in the first case, as non-klt centres are non-canonical. Comparing the discrepancies of the differentials of adjunction for Δ_k and Γ_k as in the cited proof shows that ϕ_k, f_k , and thus ϕ are birational contractions. On the other hand, if f is a Mori fibre space, then, as the dimension of the fibers of $f : X \rightarrow Y$ are upper-semicontinuous, f_k is not birational. \square

Theorem 51. [HMX2014, 3.2] *Let (X, Δ) be a \mathbb{Q} -factorial klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$ contains no non-canonical centres of (X_k, Δ_k) and is log smooth. Let $f : X \rightarrow Y$ be a step of the $(K_X + \Delta)$ -MMP. Then the minimal model program with scaling terminates using only contractions.*

Proof. (Similar proof to the cited theorem). Let $f : X \rightarrow Y$ the $(K_X + \Delta)$ -MMP with scaling of an ample divisor A . Let $\Gamma = f_*\Delta$ and $B = f_*A$. By construction, $K_Y + tB + \Gamma$ is nef for some $t > 0$. By Lemma 50, $f : X \rightarrow Y$ is a birational contraction and $f_k : X_k \rightarrow Y_k$ is a

birational contractions from $(X_k, tA_k + \Delta_k)$. If $K_X + \Delta$ is not pseudo-effective, then for $t > 0$, the result of f is a Mori fiber space, and by Lemma 50, Y_k is covered by $K_{Y_k} + tB_k + \Gamma_k$ -negative curves, contradicting bigness of $K_{X_k} + tA_k + \Delta_k$. Thus $K_X + \Delta$ is pseudo-effective, and given any $\epsilon > 0$, we may run the MMP until $t < \epsilon$. Now we conclude by Theorem 27. Letting ϵ be the constant given in that theorem, the minimal models for $(X_k, tA_k + \Delta_k)$ are all isomorphic $t \in [0, \epsilon]$. Thus, once $t < \epsilon$, any more steps in the minimal model program with scaling must be an isomorphism on the special fiber, and thus an isomorphism. Thus there exists a minimal model (Y, Γ) for (X, Δ) . \square

16. EXISTENCE OF MINIMAL MODELS GENERAL CASE

Theorem 52. *Let (X, Δ) be a klt pair of relative dimension 2, proper over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is \mathbb{Q} -Cartier, and log smooth over R . Then the minimal model of (X, Δ) exists.*

Proof. I again start by repeating the reduction of [HMX2014, 6.1]. If necessary, extend R to be complete with algebraically closed residue field k , so the strata of Δ have irreducible fibers over R . Let $f_0 : Y_0 \rightarrow X_0$ be the birational morphism of Lemma 47. Under the log smooth hypothesis, and since the strata of Δ have irreducible fibers, and f_0 blows up strata of Δ_0 , extend f_0 to a birational morphism $f : Y \rightarrow X/R$ which is a composition of smooth blow ups of strata of Δ . Write

$$K_Y + \Gamma = f^*(K_X + \Delta) + E$$

with $\Gamma \geq 0$ and $E \geq 0$, $f_*\Gamma = \Delta$, and $f_*E = 0$. Then (Y, Γ) is log smooth and the fibres of components of Γ are irreducible. By Lemma 49, (X, Δ) has a minimal model if (Y, Γ_ϵ) has a good minimal model, where $(Y, \Gamma_\epsilon) = (Y, \Gamma + \epsilon F)$ is the ϵ -log smooth model of (X, Δ) with F the sum of f -exceptional divisors and $0 < \epsilon \ll 1$.

Replace (X, Δ) by (Y, Γ_ϵ) and set $\Theta_k = \Delta_k - \Delta_k \wedge N_\sigma(X_k, K_{X_k} + \Delta_k)$ so that $\mathbb{B}_-(X_k, K_{X_k} + \Theta_k)$ contains no strata of Θ_k . Let $0 \leq \Theta \leq \Delta$ be the unique divisor such that $\Theta_k = \Theta|_{X_k}$. Let H relatively ample such that $(X, \Theta + H)$ is log smooth over R , and $K_X + \Theta + H$ is big. There is a commutative diagram:

$$\begin{array}{ccc} \pi_* \mathcal{O}_X(m(K_X + \Theta) + H) & \longrightarrow & \pi_* \mathcal{O}_X(m(K_X + \Delta) + H) \\ \downarrow & & \downarrow \\ H^0(X_k, \mathcal{O}_{X_k}(m(K_{X_k} + \Theta_k) + H_k)) & \longrightarrow & H^0(X_k, \mathcal{O}_{X_k}(m(K_{X_k} + \Delta_k) + H_k)) \end{array}$$

with surjective columns by Theorem 20 and [Liu, 5.3.20(b)], and with the bottom row an isomorphism. Applying Nakayama's lemma gives an isomorphism on the top row, so that $\Theta \geq \Delta - \Delta \wedge N_\sigma(X, K_X + \Delta)$. Again applying Theorem 20, gives $\Theta = \Delta - \Delta \wedge N_\sigma(X, K_X + \Delta)$. Thus $\Delta - \Theta \leq N_\sigma(X, K_X + \Delta)$, so by Lemma 48, it suffices to find a minimal model for (X, Θ) . Replacing (X, Δ) by (X, Θ) , it suffices to assume that $\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$ contains no strata of Δ_0 . Letting A be an ample divisor, we run the minimal model program with scaling of A . Now the assumptions of Theorem 51 apply, so we know that (X, Θ) has a minimal model. \square

REFERENCES

- [A1] Artin, M. "Algebraization of formal moduli. I, Global Analysis (Papers in Honor of K. Kodaira), 21–71." (1969).
- [A2] Artin, Michael. "Algebraization of formal moduli: II. Existence of modifications." *Annals of Mathematics* (1970): 88-135.
- [BCHM] Birkar, Caucher, et al. "Existence of minimal models for varieties of log general type." *Journal of the American Mathematical Society* 23.2 (2009): 405.
- [Bo] Bosch, Siegfried, et al. "Néron models." (1990).
- [CL] Corti, Alessio, and Vladimir Lazić. "New outlook on the minimal model program, II." *Mathematische Annalen* 356.2 (2013): 617-633.
- [CL2013] Cascini, Paolo, and Vladimir Lazić. "On the number of minimal models of a log smooth threefold." *Journal de Mathématiques Pures et Appliquées* 102.3 (2014): 597-616.
- [CP2014] Cossart, Vincent, and Olivier Piltant. "Resolution of Singularities of Arithmetical Threefolds II." arXiv preprint arXiv:1412.0868 (2014).
- [CZ] Cascini, Paolo, and DEQI ZHANG. "Effective finite generation for adjoint rings." arXiv preprint arXiv:1203.5204 (2012).
- [DF] Di Cerbo, Gabriele, and Andrea Fanelli. "Effective Matsusaka's Theorem for surfaces in characteristic p ." arXiv preprint arXiv:1501.07299 (2015).
- [EGSurfDIOP] Egbert, Andrew. SurfDIOP. <https://divisibility.wordpress.com/2015/06/12/mixed-characteristic-minimal-models-and-invariance-of-plurigenera-for-relative-dimension-2/>
- [EgSyz] Egbert, Andrew. <https://divisibility.wordpress.com/2015/08/07/adjoint-rings-and-szygies-in-mixed-or-positive-characteristic/>
- [EgDeg] Egbert, Andrew. <https://divisibility.wordpress.com/2015/08/07/adjoint-rings-and-szygies-in-mixed-or-positive-characteristic/>

- [ELM] Ein, Lawrence, et al. "Asymptotic invariants of base loci." *Annales de l'institut Fourier*. Vol. 56. No. 6. 2006.
- [Ek] Ekedahl, Torsten. "Canonical models of surfaces of general type in positive characteristic." *Publications Mathématiques de l'IHÉS* 67.1 (1988): 97-144.
- [GH] Griffiths, Phillip, and Joseph Harris. *Principles of algebraic geometry*. John Wiley & Sons, 2014.
- [GL] Gongyo, Yoshinori, and Brian Lehmann. "Reduction maps and minimal model theory." *Compositio Mathematica* 149.02 (2013): 295-308.
- [GP] Gallego, Francisco Javier, and B. P. Purnaprajna. "Projective normality and syzygies of algebraic surfaces." *Journal für die reine und angewandte Mathematik (Crelles Journal)* 1999.506 (1999): 145-180.
- [Gr] Grothendieck, A. "Éléments de géométrie algébrique." New York (1967).
- [Har] Hartshorne, Robin. *Algebraic geometry*. Vol. 52. Springer Science & Business Media, 1977.
- [H] Hashizume, Kenta. "Finite generation of adjoint ring for log surfaces." *arXiv preprint arXiv:1505.02282* (2015).
- [HMX2010] Hacon, Christopher, James McKernan, and Chenyang Xu. "On the birational automorphisms of varieties of general type." *arXiv preprint arXiv:1011.1464* (2010).
- [HMX2014] Hacon, CHRISTOPHER D., James McKernan, and Chenyang Xu. "Boundedness of moduli of varieties of general type." *arXiv preprint arXiv:1412.1186* (2014).
- [HK] Hu, Yi, and Sean Keel. "Mori dream spaces and GIT." *arXiv preprint math/0004017* (2000).

- [JKW] Elizondo, E. Javier, Kazuhiko Kurano, and Kei-ichi Watanabe. "The total coordinate ring of a normal projective variety." *Journal of Algebra* 276.2 (2004): 625-637.
- [Ke] Keel, Seán. "Basepoint freeness for nef and big line bundles in positive characteristic." *Annals of Mathematics* 149 (1999): 253-286.
- [KM] Kollár, János, and Shigefumi Mori. *Birational geometry of algebraic varieties*. Vol. 134. Cambridge University Press, 2008.
- [KMM] Kawamata, Yujiro, Katsumi Matsuda, and Kenji Matsuki. "Introduction to the minimal model problem." *Adv. Stud. Pure Math* 10 (1987): 283-360.
- [Ko2013] Kollár, János. *Singularities of the minimal model program*. Vol. 200. Cambridge University Press, 2013.
- [KU] Katsura, Toshiyuki, and Kenji Ueno. "On elliptic surfaces in characteristic p ." *Mathematische Annalen* 272.3 (1985): 291-330.
- [KK94] Kollár, János, and Sándor Kovács. "Birational geometry of log surfaces." preprint (1994).
- [Lai] Lai, Ching-Jui. "Varieties fibered by good minimal models." *Mathematische Annalen* 350.3 (2011): 533-547.
- [Laz] Lazarsfeld, Robert K. *Positivity in algebraic geometry*. Springer Science & Business Media, 2004.
- [Liu] Liu, Qing, and Reinie Erne. *Algebraic geometry and arithmetic curves*. Oxford university press, 2002.
- [MP] Manin, Yuri Ivanovic, and Alexei A. Panchishkin. *Introduction to modern number theory*. Vol. 49. Springer-Verlag Berlin Heidelberg, 2005.
- [SC] Shokurov, Vyacheslav, and Sung Choi. "Geography of log models: theory and applications." *Open Mathematics* 9.3 (2011): 489-534.
- [Suh] Suh, Junecue. "Plurigenera of general type surfaces in mixed characteristic." *Compositio Mathematica* 144.05 (2008): 1214-1226.

- [Tan] Tanaka, Hiromu. "Minimal models and abundance for positive characteristic log surfaces." *Nagoya Mathematical Journal* (2015).
- [Tan2] Tanaka, Hiromu. "The X-method for klt surfaces in positive characteristic." arXiv preprint arXiv:1202.2497 (2012).
- [Ter] Terakawa, Hiroyuki. "The d-very ampleness on a projective surface in positive characteristic." *Pacific J. of Math* 187 (1999): 187-198.
- [Xu] Xu, Chenyang. "On base point free theorem of threefolds in positive characteristic." arXiv preprint arXiv:1311.3819 (2013).