

MINIMAL MODELS FOR ARITHMETIC THREEFOLD GERMS

ANDREW EGBERT

ABSTRACT. In this note, I attempt to finish the log minimal model program proofs started in the SurfDIOP and Adjoint rings paper (see divisibility.wordpress.com). In SurfDIOP, finite generation of the canonical ring for klt, log smooth pairs on a scheme having two dimensional fibers over a DVR. In the syzygies and Adjoint rings note, the minimal model program for such (general type) pairs is studied (and the case that serre duality holds for X). In this, I relax the assumption on kodaira dimension. Sometime in the next week, I'll likely try to combine all three of these in a sane (in terms of organization) manner. Most of this paper is just using results from characteristic 0, checking quickly if they hold over a mixed characteristic DVR, and then applying the finite generation in the big case (from [\[EGSurfDIOP\]](#)).

1. FINITENESS OF MODELS FOR SURFACES

Theorem 1. *Let X be a two dimensional normal variety over a perfect field of characteristic $p > 0$. Let (X, Δ) be a KLT pair and A an ample \mathbb{Q} -divisor on X . Then there exists an $\epsilon > 0$, such that the minimal models (and the output of the minimal model program with scaling) for $(X, \Delta + tA)$ are all isomorphic for $t \in [0, \epsilon]$.*

Proof. (Essentially same as [\[Lai, 26,27\]](#) - I'm not sure if this is the correct cite for Lai's thesis which is what I am intending to cite, as it

appears to be a springerlink thingy - in any case the proof is essentially the same or simplified, but restated for the slightly different setting, and simplifies due to the fact that birational isomorphism in codimension 1 on a surface gives an isomorphism. There is also a proof of essentially the same result in [HMX2014]). Let $\phi : (X, \Delta) \rightarrow (X_g, \Delta_g)$ a good minimal model (exists by [Tan]). Let $f : X_g \rightarrow Z = \text{Proj } R(K_{X_g} + \Delta_g)$ the contraction (exists by since $K_{X_g} + \Delta_g$ is nef, hence semiample by [Tan]). Then ϕ contracts the divisorial part of $\mathbb{B}(K_X + \Delta)$. Pick $t_0 > 0$ such that $(X_g, \Delta_g + t_0\phi_*A) = (X_g, \Delta_g + t_0A_g)$ is klt with A ample on X (note A_g is big, not in general nef). For H ample on X_g , let $\phi : X_g \rightarrow X'$ be the minimal model program with scaling, which terminates again by [Tan], and gives a minimal model of $(X_g, \Delta_g + t_0\Delta_g)$ over Z . For any curve contracted by f , $(K_{X_g} + \Delta_g) \cdot C = 0$, hence $K_{X'} + \psi_*\Delta_g = K_{X'} + \Delta' \equiv_Z 0$. Thus curves contracted by ψ have trivial intersection with $K_{X_g} + \Delta_g$, and intersect negatively with A_g . Thus changing t doesn't affect which curves intersect negatively with $K_{X_g} + \Delta_g + tA_g$, and so X' is a minimal model of $(X_g, \Delta_g + tA_g)$ for all $t \in (0, t_0]$.

Now $\Delta' + t_0A'$, $A' = \psi_*A_g$ is big implies by the cone theorem (again found in [Tan]), there exist only finitely many $K_{X'} + \Delta' + t_0A'$ negative extremal rays in $\overline{NE}(X')$. These are all necessarily just intersecting A' negatively, so decreasing t_0 , eventually we get to a point where further decrease in t_0 doesn't change the number of negative extremal rays. Pick such a t_0 (Now we are on X' , not over Z). Again shrinking t_0 , suppose that $\psi \circ \phi$ is discrepancy negative w'r't $(X, \Delta + tA)$ for all t in the half open interval $(0, t_0]$. Note that $\mathbb{B}(K_X + \Delta + t_0A) \subset$

$\mathbb{B}(K_X + \Delta)$, so ψ contracts only the things not contracted by ϕ (which is no divisors). Thus $\psi \circ \phi$ is discrepancy negative on the closed interval $[0, t_0]$, so X' is a minimal model of $(X, \Delta + tA)$ for all $t \in [0, t_0]$. Thus $\mathbb{B}(K_X + \Delta + tA)$ has the same divisorial components for all $t \in [0, t_0]$. So any two minimal models for different $t \in [0, t_0]$ are isomorphic in codimension one, and as the total dimension is two, they are thus isomorphic. \square

2. CONE, RATIONALITY, AND CONTRACTION / R

(See [EGSurfDIOP] 3.12+)

3. BIG ABUNDANCE / R

(See [EGSurfDIOP, 3.12+])

4. CONTRACTIONS GIVEN BY EXTREMAL RAYS / R

(See [EGSurfDIOP, 3.12+])

5. ABUNDANT TERMINATION WITH SCALING

Theorem 2. [BCHM, 7.1] *Let $\pi : X \rightarrow U$ a projective morphism of normal quasi-projective schemes of dimension at most 3. Let V a finite dimensional affine subspace of $WDiv_{\mathbb{R}}(X)$ defined over the rationals. Fix a general ample \mathbb{Q} -divisor A over U . Let $\mathcal{C} \subset \mathcal{L}_A(V)$ a rational polytope such that if $\Delta \in \mathcal{C}$, then $K_X + \Delta$ is kawamata log terminal. Then there are finitely many rational maps $\phi_i : X \rightarrow Y_i$ over U $1 \leq i \leq k$, with the property that if $\Delta \in \mathcal{C} \cap \mathcal{E}_{A,\pi}(V)$ then there is an index $1 \leq i \leq k$ such that ϕ_i is a log terminal model of $K_X + \Delta$ over U .*

Proof. As in the source, this holds by termination in the big case (holds by [EgSyz]) and non-vanishing in the big case (see [EGSurfDIOP] 3.12+). \square

Theorem 3. *Let $(X, \Delta)/R$ be a two dimensional klt pair projective over a dvr R with a good minimal model over R . Then any $(K_X + \Delta)$ -minimal model program over R with scaling of an ample divisor terminates.*

Proof. (c.f. [Lai, 26], similar proof to the “Finiteness of Models” above). Let $\phi : (X, \Delta) \rightarrow (X_g, \Delta_g)$ a good minimal model (exists by hypothesis). Let $f : X_g \rightarrow Z = \text{Proj } R(K_{X_g} + \Delta_g)$ the contraction (exists since it’s a “Good” minimal model). Then ϕ contracts the divisorial part of $\mathbb{B}(K_X + \Delta)$. Pick $t_0 > 0$ such that $(X_g, \Delta_g + t_0\phi_*A) = (X_g, \Delta_g + t_0A_g)$ is klt with A ample on X (note A_g is big, not in general nef). For H ample on X_g , let $\psi : X_g \rightarrow X'$ be the minimal model program with scaling, which terminates by [EgSyz] (as $\Delta_g + t_0A_g$ is big), and gives a minimal model of $(X_g, \Delta_g + t_0\Delta_g)$ over Z . For any curve contracted by f , $(K_{X_g} + \Delta_g) \cdot C = 0$, hence $K_{X'} + \psi_*\Delta_g = K_{X'} + \Delta' \equiv_Z 0$. Thus curves contracted by ψ have trivial intersection with $K_{X_g} + \Delta_g$, and intersect negatively with A_g . Thus changing t doesn’t affect which curves intersect negatively with $K_{X_g} + \Delta_g + tA_g$, and so X' is a minimal model of $(X_g, \Delta_g + tA_g)$ for all $t \in (0, t_0]$.

Now $\Delta' + t_0A'$, $A' = \psi_*A_g$ is big implies by the cone theorem ([EGSurfDIOP] 3.12+), there exist only finitely many $K_{X'} + \Delta' + t_0A'$ negative extremal rays in $\overline{NE}(X')$. These are all necessarily just

intersecting A' negatively, so decreasing t_0 , eventually we get to a point where further decrease in t_0 doesn't change the number of negative extremal rays. Pick such a t_0 (Now we are on X' , not over Z). Again shrinking t_0 , suppose that $\psi \circ \phi$ is discrepancy negative w'r't $(X, \Delta + tA)$ for all t in the half open interval $(0, t_0]$. Note that $\mathbb{B}(K_X + \Delta + t_0A) \subset \mathbb{B}(K_X + \Delta)$, so ψ contracts only the things not contracted by ϕ (which is no divisors). Thus $\psi \circ \phi$ is discrepancy negative on the closed interval $[0, t_0]$, so X' is a minimal model of $(X, \Delta + tA)$ for all $t \in [0, t_0]$. Thus $\mathbb{B}(K_X + \Delta + tA)$ has the same divisorial components for all $t \in [0, t_0]$. So any two minimal models for different $t \in [0, t_0]$ are isomorphic in codimension one.

The rest holds as in the cited proof, with finiteness of models by Theorem 2. □

6. REDUCTIONS

Theorem 4. [CP2014, 1.1] *Let X be a reduced and separated Noetherian scheme which is quasi-excellent and of dimension at most three. There exists a proper birational morphism $\pi : X' \rightarrow X$ with the following properties:*

- (i) X' is everywhere regular
- (ii) π induces an isomorphism $\pi^{-1}(\text{Reg } X) \approx \text{Reg } X$
- (iii) $\pi^{-1}(\text{Sing } X)$ is a strict normal crossings divisor on X' .

Lemma 5. [HMX2014, 2.8.3] *Let (X, Δ) be a log smooth pair which is a scheme of dimension at most 3 (or wherever resolution holds) where*

the coefficients of Δ belong to $(0, 1]$ and X is projective. If (X, Δ) has a weak log canonical model, then there is a sequence $\pi : Y \rightarrow X$ of smooth blow ups of the strata of Δ such that if we write $K_Y + \Gamma = \pi^*(K_X + \Delta) + E$, where $\Gamma \geq 0$ and $E \geq 0$ have no common components, $\pi_*\Gamma = \Delta$ and $\pi_*E = 0$ and if we write

$$\Gamma' = \Gamma - \Gamma \wedge N_\sigma(Y, K_Y + \Gamma),$$

then $\mathbb{B}_-(Y, K_Y + \Gamma')$ contains no strata of Γ' . If Δ is a \mathbb{Q} -divisor, then Γ' is a \mathbb{Q} -divisor.

Proof. (same as source which is stated over \mathbb{C} , since just using resolution). □

Lemma 6. [HX, 2.10] *Let $f : X \rightarrow U$ a projective morphism from a scheme of dimension at most 3, to a scheme of dimension at most 3, (X, Δ) a dlt pair and $\mu : X' \rightarrow X$ a proper birational morphism. Suppose that termination is satisfied for general type scheme pairs (X, Φ) of the same dimension. Write $K_{X'} + \Delta' = \mu^*(K_X + \Delta) + F$ where Δ' and F are effective with no common components. Then (X, Δ) has a good minimal model over U if and only if (X', Δ') has a good minimal model over U .*

Proof. Same as in the source, noting that resolution holds in this dimension, and the termination for the big case holds by [EgSyz], and termination in the abundant case holds by Theorem 3. □

Lemma 7. [HMX2014, 5.3] *Let (X, Δ) be a dlt pair with X \mathbb{Q} -factorial and projective and Δ a \mathbb{Q} -divisor. If Φ is a \mathbb{Q} -divisor such that*

$$0 \leq \Delta - \Phi \leq N_\sigma(X, K_X + \Delta),$$

then (X, Φ) has a good minimal model implies (X, Δ) has a good minimal model.

Proof. This essentially follows as in the source, applying the Cone Theorem in [EGSurfDIOP, 3.12+] as needed. (I'll expand on this more, since it's an lengthy reduction, but seems to use only the Cone Theorem). \square

Lemma 8.

7. MAIN THEOREM

Theorem 9. *Let (X, Δ) be a klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is \mathbb{Q} -Cartier, pseudo-effective, and log smooth over R . Then the minimal model of (X, Δ) exists.*

Proof. I start by repeating the reduction of [HMX2014, 6.1]. If necessary, extend R to be complete with algebraically closed residue field k , so the strata of Δ have irreducible fibers over R . Let $f_0 : Y_0 \rightarrow X_0$ be the birational morphism of Lemma 5. Under the log smooth hypothesis, and since the strata of Δ have irreducible fibers, and f_0 blows up strata of Δ_0 , extend f_0 to a birational morphism $f : Y \rightarrow X/R$ which

is a composition of smooth blow ups of strata of Δ . Write

$$K_Y + \Gamma = f^*(K_X + \Delta) + E$$

with $\Gamma \geq 0$ and $E \geq 0$, $f_*\Gamma = \Delta$, and $f_*E = 0$. (Y, Γ) is log smooth and the fibres of components of Γ are irreducible. As (X_0, Δ_0) is projective variety of dimension 2, by [Tan] it has a good minimal model, so (Y_0, Δ_0) has a good minimal model by Lemma 6, and similarly (X, Δ) has a good minimal model iff (Y, Γ) has a good minimal model. Thus it suffices to show that (Y, Γ) has a good minimal model. Now we are in close to the situation of the main invariance of plurigenera theorem of [EGSurfDIOP], but with possibly slightly worse singularities.

Replace (X, Δ) by (Y, Γ) and set $\Theta_0 = \Delta_0 - \Delta_0 \wedge N_\sigma(X_0, K_{X_0} + \Delta_0)$ so that $\mathbb{B}_-(X_0, K_{X_0})$ contains no strata of Θ_0 . Let $0 \leq \Theta \leq \Delta$ be the unique divisor such that $\Theta_0 = \Theta|_{X_0}$. Let H relatively ample such that $(X, \Delta + H)$ is log smooth over R , and $kK_X + H$ is big. There is a commutative diagram:

$$\begin{array}{ccc} \pi_*\mathcal{O}_X(m(K_X + \Theta) + H) & \longrightarrow & \pi_*\mathcal{O}_X(m(K_X + \Delta) + H) \\ \downarrow & & \downarrow \\ H^0(X_0, \mathcal{O}_{X_0}(m(K_{X_0} + \Theta_0) + H_0)) & \longrightarrow & H^0(X_0, \mathcal{O}_{X_0}(m(K_{X_0} + \Delta_0) + H_0)) \end{array}$$

with surjective columns by the main invariance of plurigenera theorem of [EGSurfDIOP], with the bottom row an isomorphism. Applying Nakayama's lemma as in [EGSurfDIOP, EgSyz] gives an isomorphism on the top row, so that $\Theta \geq \Delta - \Delta \wedge N_\sigma(X, K_X + \Delta)$. Again applying the invariance of plurigenera in [EGSurfDIOP], gives $\Theta = \Delta - \Delta \wedge$

$N_\sigma(X, K_X + \Delta)$. Thus $\Delta - \Theta \leq N_\sigma(X, K_X + \Delta)$, so by Lemma 7, it suffices to find a minimal model for (X, Θ) . Replace (X, Δ) by (X, Θ) , it suffices to assume that $\mathbb{B}_-(X_0, K_{X_0} + \Delta_0)$ contains no strata of Δ_0 .

Let A be an ample divisor, we run the minimal model program with scaling of A . Now the assumptions of [HMX2014, 3.1] apply, so we can actually find each step $f : X^i \rightarrow Y^i$ of the minimal model program with scaling induces a (WLCM) $f_0 : X_0^i \rightarrow Y_0^i$ of $(X_0, tA_0 + \Delta_0)$. Now applying finiteness of models gives the termination. Now, we know that (X, Φ) has a minimal model, (X', Φ') with $K_{X'} + \Phi'$ semiample on both fibers (by [Tan]). Now for any $\epsilon > 0$, and any geometric valuation Γ on X' , $\sigma_\Gamma(K_{X'_0} + \Phi'_0) = \sigma_\Gamma(K_{X'_\eta} + \Phi'_\eta) = 0$ so that σ_Γ is identically zero, and thus the centre of Γ is not in $\mathbb{B}(K_{X'} + \Phi')$ for any such Γ . \square

REFERENCES

- [A1] Artin, M. "Algebraization of formal moduli. I, Global Analysis (Papers in Honor of K. Kodaira), 21–71." (1969).
- [A2] Artin, Michael. "Algebraization of formal moduli: II. Existence of modifications." *Annals of Mathematics* (1970): 88-135.
- [BC] Baker, Matthew., and Janos A. Csirik. "On the Isomorphism Between the Dualizing Sheaf and the Canonical Sheaf." (1996).
- [BCHM] Birkar, Caucher, et al. "Existence of minimal models for varieties of log general type." *Journal of the American Mathematical Society* 23.2 (2009): 405.
- [Bir] Birkar, Caucher. "Existence of flips and minimal models for 3-folds in char p." arXiv preprint arXiv:1311.3098 (2013).
- [Bo] Bosch, Siegfried, et al. "Néron models." (1990).

- [BP] Berndtsson, Bo, and Mihai Paun. "Quantitative extensions of pluricanonical forms and closed positive currents." *Nagoya Math. J* 205 (2012): 25-65.
- [CL] Corti, Alessio, and Vladimir Lazić. "New outlook on the minimal model program, II." *Mathematische Annalen* 356.2 (2013): 617-633.
- [CP2009] Cossart, Vincent, and Olivier Piltant. "Resolution of singularities of threefolds in positive characteristic II." *Journal of Algebra* 321.7 (2009): 1836-1976.
- [CP2014] Cossart, Vincent, and Olivier Piltant. "Resolution of Singularities of Arithmetical Threefolds II." arXiv preprint arXiv:1412.0868 (2014).
- [CR] Chatzistamatiou, Andre, and Kay Rülling. "Higher direct images of the structure sheaf in positive characteristic." *Algebra & Number Theory* 5.6 (2012): 693-775.
- [CZ] Cascini, Paolo, and DEQI ZHANG. "Effective finite generation for adjoint rings." arXiv preprint arXiv:1203.5204 (2012).
- [Das] Das, Omprokash. "On Strongly F -Regular Inversion of Adjunction." arXiv preprint arXiv:1310.8252 (2013).
- [Deb] Dèbes, Pierre, et al., eds. *Arithmetic and geometry around Galois theory*. Springer Science & Business Media, 2012.
- [DF] Di Cerbo, Gabriele, and Andrea Fanelli. "Effective Matsusaka's Theorem for surfaces in characteristic p ." arXiv preprint arXiv:1501.07299 (2015).
- [DI] Deligne, Pierre, and Luc Illusie. "Relèvements modulop 2 et décomposition du complexe de de Rham." *Inventiones Mathematicae* 89.2 (1987): 247-270.
- [EGSurfDIOP] Egbert, Andrew. SurfDIOP. <https://divisibility.wordpress.com/2015/06/12/mixed-characteristic-minimal-models-and-invariance-of-plurigenera-for-relative-dimension-2/>

- [EgSyz] Egbert, Andrew. <https://divisibility.wordpress.com/2015/08/07/adjoint-rings-and-szygies-in-mixed-or-positive-characteristic/>
- [ELM] Ein, Lawrence, et al. "Asymptotic invariants of base loci." *Annales de l'institut Fourier*. Vol. 56. No. 6. 2006.
- [Ek] Ekedahl, Torsten. "Canonical models of surfaces of general type in positive characteristic." *Publications Mathématiques de l'IHÉS* 67.1 (1988): 97-144.
- [EV] Viehweg, Eckart. *Lectures on vanishing theorems*. Vol. 20. Springer Science & Business Media, 1992.
- [GH] Griffiths, Phillip, and Joseph Harris. *Principles of algebraic geometry*. John Wiley & Sons, 2014.
- [Gra] Grauert, Hans, Thomas Peternell, and Reinhold Remmert. *Several complex variables VII: sheaf-theoretical methods in complex analysis*. Vol. 7. Springer Science & Business Media, 1994.
- [Gr] Grothendieck, A. "Éléments de géométrie algébrique." New York (1967).
- [GW] Görtz, Ulrich, and Torsten Wedhorn. *Algebraic Geometry*. Vieweg+Teubner, 2010.
- [HK] Hacon, Christopher D., and Sándor Kovács. *Classification of higher dimensional algebraic varieties*. Vol. 41. Springer Science & Business Media, 2011.
- [Hara98] Hara, Nobuo. "Classification of two-dimensional F-regular and F-pure singularities." *Advances in Mathematics* 133.1 (1998): 33-53.
- [Har] Hartshorne, Robin. *Algebraic geometry*. Vol. 52. Springer Science & Business Media, 1977.
- [HS] Hindry, Marc, and Joseph H. Silverman. *Diophantine geometry: an introduction*. Vol. 201. Springer Science & Business Media, 2000.

- [HMX2010] Hacon, Christopher, James McKernan, and Chenyang Xu. "On the birational automorphisms of varieties of general type." arXiv preprint arXiv:1011.1464 (2010).
- [HMX2014] Hacon, CHRISTOPHER D., James McKernan, and Chenyang Xu. "Boundedness of moduli of varieties of general type." arXiv preprint arXiv:1412.1186 (2014).
- [HX] Hacon, Christopher D., and Chenyang Xu. "Existence of log canonical closures." *Inventiones mathematicae* 192.1 (2013): 161-195.
- [KU] Katsura, Toshiyuki, and Kenji Ueno. "On elliptic surfaces in characteristic." *Mathematische Annalen* 272.3 (1985): 291-330.
- [KK94] Kollár, János, and Sándor Kovács. "Birational geometry of log surfaces." preprint (1994).
- [KM] Kollár, János, and Shigefumi Mori. *Birational geometry of algebraic varieties*. Vol. 134. Cambridge University Press, 2008.
- [Ko] Kollár, János. *Singularities of the minimal model program*. Vol. 200. Cambridge University Press, 2013.
- [Ko2] Kollár, János. *Shafarevich maps and automorphic forms*. Princeton University Press, 2014.
- [Ko1991] Kollár, János, ed. *Flips and abundance for algebraic threefolds: a summer seminar at the University of Utah, Salt Lake City, 1991*. Société mathématique de France, 1992.
- [Lai] Lai, Ching-Jui. "Varieties fibered by good minimal models." *Mathematische Annalen* 350.3 (2011): 533-547.
- [Laz] Lazarsfeld, Robert K. *Positivity in algebraic geometry*. Springer Science & Business Media, 2004.
- [Lev] Levine, Marc. "Pluri-canonical divisors on Kähler manifolds." *Inventiones mathematicae* 74.2 (1983): 293-303.
- [Lev-2] Levine, Marc. "Pluri-canonical divisors on Kähler manifolds, II." *Duke Math. J* 52.1 (1985): 61-65.

- [Liu] Liu, Qing, and Reinie Erne. Algebraic geometry and arithmetic curves. Oxford university press, 2002.
- [LS] Liedtke, Christian, and Matthew Satriano. "On the birational nature of lifting." *Advances in Mathematics* 254 (2014): 118-137.
- [Ma] Maddock, Zachary. "A bound on embedding dimensions of geometric generic fibers." arXiv preprint arXiv:1407.2529 (2014).
- [Mum] Mumford, David, Chidambaram Padmanabhan Ramanujam, and Jurij Ivanovič Manin. Abelian varieties. Vol. 48. Oxford: Oxford university press, 1970.
- [Oss] Osserman, Brian. "Notes on Cohomology and Base Change." <https://www.math.ucdavis.edu/~osserman/math/cohom-base-change.pdf>
- [Sat] Satriano, Matthew. "De Rham theory for tame stacks and schemes with linearly reductive singularities." arXiv preprint arXiv:0911.2056 (2009).
- [Serre] Serre, Jean-Pierre. Local fields. Vol. 67. Springer Science & Business Media, 2013.
- [Siu] Siu, Yum-Tong. "Invariance of plurigenera." arXiv preprint alg-geom/9712016 (1997).
- [Siu2] Siu, Yum-Tong. "Extension of twisted pluricanonical sections with plurisubharmonic weight and invariance of semipositively twisted plurigenera for manifolds not necessarily of general type." *Complex geometry*. Springer Berlin Heidelberg, 2002. 223-277.
- [Suh] Suh, Junecue. "Plurigenera of general type surfaces in mixed characteristic." *Compositio Mathematica* 144.05 (2008): 1214-1226.
- [Tan] Tanaka, Hiromu. "Minimal models and abundance for positive characteristic log surfaces." *Nagoya Mathematical Journal* (2015).
- [Tan2] Tanaka, Hiromu. "The X-method for klt surfaces in positive characteristic." arXiv preprint arXiv:1202.2497 (2012).

- [Tan3] Tanaka, Hiromu. "The trace map of Frobenius and extending sections for threefolds." arXiv preprint arXiv:1302.3134 (2013).
- [Tan4] Tanaka, Hiromu. "Minimal model theory for surfaces over an imperfect field." arXiv preprint arXiv:1502.01383 (2015).
- [Ter] Terakawa, Hiroyuki. "The d -very ampleness on a projective surface in positive characteristic." *Pacific J. of Math* 187 (1999): 187-198.
- [Wal] Waldron, Joe. "Finite generation of the log canonical ring for 3-folds in char p ." arXiv preprint arXiv:1503.03831 (2015).