

MINIMAL MODELS FOR ARITHMETIC THREEFOLDS

ABSTRACT. I study the existence of log minimal models for a pair of relative dimension 2 over a Dedekind Domain . This generalizes the semistable result of Kawamata [Kaw94, Kaw98]. Also I note the invariance of log plurigenera for such pairs, generalizing the result of [Suh08]. To extend the result from DVR's to Dedekind Domains, some computability results (which may be interesting by themselves) are given for basepoint-freeness, Kawamata-Viehweg vanishing, and finite generation of klt log pairs on the positive characteristic surface.

CONTENTS

1. Introduction	2
Part 1. Local Minimal Model Program	3
2. Intersection Theory	3
3. Log Resolution	5
4. KLT Kawamata Viehweg Vanishing	6
5. Invariance of Plurigenera Special Case	7
6. Invariance of Plurigenera General Case	12
7. Finite Generation (Big Case)	13
8. Finiteness of Models For Surfaces	15
9. Big Abundance / \mathbb{R}	17
10. Cone, Rationality, and contraction / \mathbb{R}	20
11. Negativity Lemma	23

12. Flips	24
13. Running the Minimal Model Program for General Type	25
14. Reductions To General Case	28
15. Existence of Minimal Models (for germs) Special Case	30
16. Existence of Minimal Models (for germs) General Case	32
Part 2. Globalization	34
17. KLT Vanishing	34
18. Generalization of Terakawa's Basepoint Free Theorem	39
19. Minimal Models for Arithmetic Threefolds	41
References	43

1. INTRODUCTION

In this paper I prove the following Theorem:

Theorem 1. *Let (X, Δ) be a pseudoeffective klt pair of relative dimension 2 over a Dedekind Domain A . Assume that $K_X + \Delta$ is \mathbb{Q} -Cartier, and log smooth over A . Then the minimal model of (X, Δ) exists.*

The above result is somewhat related to a theorem of Kawamata [[Kaw94](#), [Kaw98](#)] where he proves the terminal semistable case of the above assuming no boundary and in characteristics $p \geq 5$. However, the techniques used in this paper are mostly quite different from Kawamata's paper, and the motivating terminal case for the above theorem would more accurately be the proof of terminal minimal models over a

DVR given by Katsura and Ueno [KU85]. Other somewhat similar results are the recent proofs of the existence of terminal and klt minimal models respectively in positive characteristic for threefolds over a field in [HX13, Bir15]. However, the case considered in this article is of geometric dimension 2 and arithmetic dimension 1, rather than geometric dimension 3 and arithmetic dimension 0. Another somewhat similar result in geometric dimension 3, arithmetic dimension 0 and characteristic $p > 5$ is [BCZ15] where the existence of log minimal models over a curve in equal characteristic is proven (note that the theorem 1 can apply in equal characteristic over a field as well even without the restriction that $p > 5$).

Theorem 1 is the result of applying the techniques of [CL13] (in the big case) and [HMX14] (in general) after achieving a generalization of Suh's Theorem on the invariance of plurigena [Suh08] (namely to Kawamata Log Terminal Pairs). Finally the result is globalized to a Dedekind Domain in a similar manner to [Kaw94]. Thanks to my advisor, Professor Hacon for help in debugging some of the proofs and for helpful suggestions.

Part 1. Local Minimal Model Program

2. INTERSECTION THEORY

Suppose X/R is a proper algebraic space of relative dimension 2 over a discrete valuation ring R with residue field k of characteristic $p > 0$ and perfect fraction field K (See for example the compact Shimura Varieties mentioned in [Suh08, 0.1]).

Lemma 2. [KU85, 9.3] *Let $\varphi : X \rightarrow \text{Spec}(R)$ be an algebraic two-dimensional space, smooth, proper, separated, and of finite type over $\text{spec}(R)$, where R is a DVR with algebraically closed residue field k , and field of fractions K . Let X_K (resp. X_k) denote the generic geometric (resp. closed) fibre of φ . If D, D' are divisors on X , then*

$$\begin{aligned} K_{X_k}^2 &= K_{X_K}^2 \\ (K_{X_k} \cdot D_k)_{X_k} &= (K_{X_K} \cdot D_K)_{X_K} \\ (D_k \cdot D'_k)_{X_k} &= (D_K \cdot D'_K)_{X_K}. \end{aligned}$$

If X/R is smooth, then Lemma 2 applies to show that for any divisors C, D extending to both fibers, we can define $C.D = (C|_{X_k} \cdot D|_{X_k})_{X_k}$. On the other hand, if X/R is merely normal and proper, but is actually a scheme, the resolution of singularities, Theorem 4, holds. Thus the intersection theory can be defined as in [T⁺14, Def 3.1]: $f : X' \rightarrow X$ is a resolution, and $C.D = f^*C \cdot f^*D$ for two divisors C, D extending to both fibers of Y/R . By properness, this intersection extends by linearity to Weil divisors with \mathbb{Q} or \mathbb{R} coefficients. Numerical equivalence and $N^1(X)_{\mathbb{Q}, \mathbb{R}}$ are then defined as usual.

Note the two different hypothesis here: invariance of the plurigenera requires a proper, smooth algebraic space of relative dimension 2, while the existence of minimal models requires the intermediate steps of the minimal model program to have some intersection theory, and thus be schemes. Recall that a Cartier divisor C on a variety is usually called nef if $C.D \geq 0$ for all curves C .

Lemma 3. *Let S be an affine Dedekind scheme and $f : X \rightarrow S$ a projective morphism. Let \mathcal{L} be an invertible sheaf on X such that \mathcal{L}_s is nef for every closed point $s \in S$. Then \mathcal{L} is nef.*

Proof. By [LE02, 5.3.24], the statement holds when nef is replaced by ample. Now, c.f. [Laz04, 1.4.10], it suffices to choose an ample divisor H which restricts to X_s , so that $D_s + \epsilon H_s$ is ample for all sufficiently small ϵ . Then $D + \epsilon H$ is ample for all sufficiently small ϵ , and so D is nef. \square

3. LOG RESOLUTION

The following statement of resolution will be used. So for example, if X is an algebraic variety over a number field K then, by [MP05, III.2], there is a completed arithmetical variety \overline{X} of dimension $\dim(X) + 1$:

$$\overline{X} \rightarrow \overline{\text{Spec } \mathcal{O}_K}.$$

(Note that a complete DVR is excellent).

Theorem 4. [CP14, 1.1] *Let X be a reduced and separated Noetherian scheme which is quasi-excellent and of dimension at most three. There exists a proper birational morphism $\pi : X' \rightarrow X$ with the following properties:*

- (i) X' is everywhere regular
- (ii) π induces an isomorphism $\pi^{-1}(\text{Reg } X) \approx \text{Reg } X'$
- (iii) $\pi^{-1}(\text{Sing } X)$ is a strict normal crossings divisor on X' .

The following Statement of Log Resolution will be used in the Log Smooth case (i.e. the case with (X, Δ) a pair with X smooth and Δ simple normal crossings.)

Theorem 5. [CP14, 4.3] *Let S be a regular Noetherian irreducible scheme of dimension three which is excellent and $\mathcal{I} \subset \mathcal{O}_S$ be a nonzero ideal sheaf. There exists a finite sequence*

$$S =: S(0) \leftarrow S(1) \leftarrow \cdots \leftarrow S(r)$$

with the following properties:

(i) for each j , $0 \leq j \leq r-1$, $S(j+1)$ is the blowing up along a regular integral subscheme $\mathcal{Y}(j) \subset \mathcal{S}(j)$ with

$$\mathcal{Y}(j) \subseteq \{s_j \in \mathcal{S}(j) : \mathcal{I}\mathcal{O}_{S(j), s_j} \text{ is not locally principal}\}.$$

(ii) $\mathcal{I}\mathcal{O}_{S(r)}$ is locally principal.

4. KLT KAWAMATA VIEHWEG VANISHING

Theorem 6. [Tan12, 2.11] *Let (X, Δ) be a normal projective klt surface over an algebraically closed field of characteristic $p > 0$, where $K_X + \Delta$ is an effective \mathbb{R} -divisor. Let N be a nef and big \mathbb{R} -cartier \mathbb{R} -divisor. Let D be a \mathbb{Q} -Cartier \mathbb{Z} -divisor such that $D - (K_X - \Delta)$ is nef and big. Then there exists an r_0 such that*

$$H^i(X, \mathcal{O}_X(D + rN + N')) = 0$$

for every $i > 0$, every positive real number $r \geq r_0$, and every nef \mathbb{R} -Cartier \mathbb{R} -divisor N' such that $rN + N'$ is a Cartier divisor.

Remark 7. Using Terakawa's Theorem [Ter99], (and possibly restricting the characteristic to be greater than 2) it is possible to prove a version of Theorem 49 where the constant r_0 is computable. A sketch appears towards the end of this article.

5. INVARIANCE OF PLURIGENERA SPECIAL CASE

Lemma 8. [KU85, 9.4] *Let $\varphi : X \rightarrow \text{Spec}(R)$ be an algebraic two-dimensional space, proper, separated, and of finite type over $\text{spec}(R)$, where R is a DVR with algebraically closed residue field k , and field of fractions K . Let X_K (resp. X_k) denote the generic geometric (resp. closed) fibre of φ . If X_k contains an exceptional curve of the first kind e , there exists a DVR $\tilde{R} \supset R$, with residue field isomorphic to k , and a proper smooth morphism $\tilde{\varphi} : \tilde{X} \rightarrow \text{Spec}(\tilde{R})$ of algebraic spaces which is separated and of finite type and a proper surjective morphism $\pi : X \otimes \tilde{R} \rightarrow \tilde{Y}$ over $\text{Spec}(\tilde{R})$ such that on the closed fibre, π induces the contraction of the exceptional curve e . Moreover, on the generic fibre, π also induces a contraction of an exceptional curve of the first kind.*

Proof. (Follows easily from the cited theorem, although I make a small observation about the extension of residue fields). By [Art69, Cor 6.2] $\text{Hilb}_{X/\text{Spec}(R)}$ is represented by an algebraic space \mathcal{H} which is locally of finite type over $\text{Spec}(R)$. Let Y be the irreducible component containing the point $\{e\}$ corresponding to the exceptional curve e on the special fiber. Then $e \approx \mathbb{P}_k^1$ and $N_{e/X} \approx \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}$, so Y is regular at $\{e\}$ and of dimension 1. Since e is fixed in the special fiber, the structure

morphism $p : Y \rightarrow \text{Spec}(R)$ is surjective. By [BLR12, 2.2.14]¹, we can find an étale cover $\tilde{R} \supset R$ and a morphism $j : \text{Spec}(\tilde{R}) \rightarrow Y$ over $\text{Spec}(R)$ with $j(\tilde{o}) = \{e\}$ (if R is not already complete, then first extend R to a complete DVR using [Gro67, 0.6.8.2,3] so that \tilde{R} is again a DVR). As k is assumed algebraically closed, and $\tilde{R} \rightarrow R$ is unramified, then the extension of residue fields is finite and separable at the closed point of R , and hence an isomorphism of residue fields. Let $\hat{p} : \mathcal{E} \rightarrow \text{Spec}(\tilde{R})$ be the pull-back of the universal family over Y . As the closed fibre \mathcal{E}_0 is a projective line, we may choose the morphism j in such a way that the generic fibre \mathcal{E}_1 of \hat{p} is also a projective line. Moreover, \mathcal{E} can be considered as a smooth closed algebraic subspace of codimension 1 in $\hat{X} = X \otimes \tilde{R}$. By Lemma 2, $-1 = \mathcal{E}_0^2 = \mathcal{E}_1^2$. Hence \mathcal{E}_1 is also an exceptional curve of the first kind. Hence by [Art70, Cor 6.11] there exists a contraction morphism $\pi : \hat{X} \rightarrow \tilde{X}$ over $\text{Spec}(\tilde{R})$ which contracts \mathcal{E} to a section of $\tilde{\varphi} : \tilde{X} \rightarrow \text{Spec}(R)$ where $\tilde{\varphi}$ is proper, smooth, separated and of finite type over $\text{Spec}(\tilde{R})$. \square

Lemma 9. [KK94, 2.3.5] *Let (S, B) be a log-canonical surface over an algebraically closed field of characteristic $p > 0$. If $C \subset S$ is a curve with $C^2 < 0$ and $C \cdot (K_S + B) < 0$, then $C \approx \mathbb{P}^1$ and it can be contracted to a log-canonical point.*

Theorem 10. [T⁺14, 4.4, 4.6] *Let X be a projective normal surface and let Δ be an effective \mathbb{R} -divisor such that $K_X + \Delta$ is \mathbb{R} -cartier. Let*

¹If k is perfect, the proof of the cited lemma seems to go through when $f : X \rightarrow S$ is merely smooth at $x \in X$ and $f : X \rightarrow S$ is locally of finite type.

$\Delta = \sum b_i B_i$ be the prime decomposition. Let H be an \mathbb{R} -cartier ample \mathbb{R} -divisor. Then the following assertions hold:

- (1) $\overline{NE}(X) = \overline{NE}(X)_{K_X + \Delta \geq 0} + \sum \mathbb{R}_{\geq 0} [C_i]$
- (2) $\overline{NE}(X) = \overline{NE}(X)_{K_X + \Delta + H \geq 0} + \sum_{finite} \mathbb{R}_{\geq 0} [C_i]$
- (3) Each C_i in (1) and (2) is rational or $C_i = B_j$ for some B_j with $B_j^2 < 0$.
- (4) Each C_i in (1) and (2) satisfies $0 < -C_i \cdot (K_X + \Delta) \leq 3$.

Theorem 11. [KK94, 2.3.6 (Log MMP For Surfaces)] *Let (S, B) be a log canonical surface (over an algebraically closed field). There exists a sequence of contractions $f : S \rightarrow S_1 \rightarrow \cdots \rightarrow S_n = S'$ such that S' is log canonical (even plt at every point where f^{-1} is not an isomorphism) and satisfies exactly one of the following conditions:*

- $K_{S'} + f_* B$ is nef
- There exists a $g : S' \rightarrow T$ morphism, such that S' is a birationally ruled surface over the curve T .
- $(S', f_* B)$ is a log Del Pezzo surface.

Remark 12. (Facts about terminal pairs) If $\dim X = 2$ and (X, D) is a terminal pair, then X is smooth [Kol13, 2.30]. By [KM08, 3.43], terminal singularities are preserved after birational contraction of a $K_X + D$ negative extremal ray, as long as the ray is not a component of D .

Proposition 13. *Let (X, Δ) be a terminal pair of relative dimension 2 over a DVR R with algebraically closed residue field k of characteristic*

$p > 0$ and perfect fraction field K . Assume that $\mathbb{B}_-(K_X + \Delta) \wedge \Delta = \emptyset$ and that $K_X + \Delta$ is \mathbb{Q} -Cartier with $\kappa(K_{X_k} + \Delta_k) = 2$. Then there exists an m_0 such that for $m_0 | m$,

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

The proof of proposition 13 is given in the following claims.

Claim. Assumptions as above, after passing to an extension R' of R , there is a proper, smooth algebraic space X^{min}/R' and an R' morphism $X \otimes R' \rightarrow X^{min}$ such that both $X_k \rightarrow X_k^{min}$ is obtained by successive blow-downs of (-1) curves and $K_{X_k^{min}} + \Delta_{X_k^{min}}$ is nef.

Proof. As k is algebraically closed of characteristic $p > 0$, then by the Cone Theorem 10,

$$\overline{NE}(X_k) = \overline{NE}(X_k)_{K_{X_k} + \Delta_k \geq 0} + \sum \mathbb{R}_{\geq 0} [C_i]$$

with each C_i is rational or $C_i = B_j$ for some B_j a component of Δ with $B_j^2 < 0$. Under the assumption that $\kappa(K_{X_k} + \Delta_k) = 2$, by Theorem 11 then it must be the case that $C_i^2 < 0$. Thus Theorem 9 implies $C_i \approx \mathbb{P}^1$ and C_i can be contracted to a log-canonical point. By Remark 12 (which applies since $\mathbb{B}_-(K_X + \Delta) \wedge \Delta = \emptyset$) the resulting pair is again terminal, hence smooth, and so it must be the case that C_i was an exceptional curve of the first kind. Therefore it is possible to apply Lemma 8.

By Lemma 8, there is a DVR $\tilde{R} \supset R$ such that $\pi : X \otimes \tilde{R} \rightarrow \tilde{X}$ induces the contraction of C_i on X_k and X_K , and further $\tilde{\varphi} : \tilde{X} \rightarrow$

$\text{Spec}(\tilde{R})$ is proper, smooth, separated, and finite type. Note that after a base change, the extension $\tilde{R} \supset R$ induces a finite extension on residue fields, and since k is algebraically closed, it induces identity on residue fields. Now I need to work with \tilde{X} . \tilde{X}_k is projective, so the same process can be repeated. Each extension $\tilde{R} \supset R$ induces a flat base change on the generic fiber and an isomorphism on the special fiber, and blow-ups are preserved by flat base-change, so there is no problem in extending R . As the Picard number of the special fiber drops each step, there are only finitely many steps. \square

Claim 14. We also have $K_{X_K^{min}} + \Delta_{X_K^{min}}$ nef.

Proof. It suffices to apply Lemma 3 to X_k^{min} , and then restrict to X_K . \square

Claim 15. There is an m_0 such that for $m_0|m$, we have

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

Proof. Suppose first that (X_k, Δ_k) is big. By the claims 5 and 14, we achieve X^{min} such that $K_X + \Delta$ is nef on both X_k^{min} and X_K^{min} . By Theorem 6 applied to the special fiber (and the semicontinuity theorem) there exists an $m_0 \gg 0$ such that for $m_0|m$ and $i > 0$, we have

$$h^i(m(K_{X_K} + \Delta_K)) = h^i(m(K_{X_k} + \Delta_k)) = 0.$$

Thus, applying the invariance of Euler characteristic [LE02, 5.3.22], and birational invariance of the plurigenera, it follows that for $m_0|m$,

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

□

Remark 16. In fact, if $\Delta = 0$, then, at the end of the above proof, Ekedahl's vanishing Theorem [Eke88] can be applied and $m_0 = 2$, in general m_0 can be computed using the computable version of Theorem 49 found later in this article.

6. INVARIANCE OF PLURIGENERA GENERAL CASE

Theorem 17. *Let (X, Δ) be a log smooth klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is big, and \mathbb{Q} -Cartier. Then, there exists an m_0 such that for $m \in m_0\mathbb{Z}^+$,*

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

Proof. Since the hypothesis and conclusion are preserved by base change, then after extending R if necessary, we may assume that k is algebraically closed and that R is complete. To begin, I make the reduction of [HMX10, 1.6]. As (X, Δ) is log smooth, then replacing (X, Δ) by a blow-up of strata, assume (X, Δ) is terminal. Recall that the log-canonical ring of $K_{X_k} + \Delta_k$ is finitely generated (c.f. [T⁺14, 7.1]) so

$N_\sigma(K_{X_k} + \Delta_k)$ is a \mathbb{Q} -divisor. Let

$$\Theta_k = \Delta_k - \Delta_k \wedge N_\sigma(K_{X_k} + \Delta_k)$$

and let $0 \leq \Theta \leq \Delta$ be a \mathbb{Q} -divisor on X/R such that, by log-smoothness $\Theta|_{X_k} = \Theta_k$. Letting $m \gg 0$ to fit the hypothesis of Proposition 13, and sufficiently divisible such that $m(K_{X_k} + \Theta_k)$ is integral, then by definition of N_σ ,

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_k} + \Theta_k)).$$

Furthermore, by Proposition 13,

$$h^0(m(K_{X_k} + \Theta_k)) = h^0(m(K_{X_K} + \Theta_K)).$$

As $\Theta \leq \Delta$, the theorem follows by semicontinuity. \square

7. FINITE GENERATION (BIG CASE)

Corollary 18. *Situation as above, after possibly extending R , the canonical ring $R(K_X + \Delta)$ is finitely generated over R .*

Proof. By [T⁺14], there is $m \gg 0$, such that

$$\bigoplus_{m \geq 0} H^0(ml(K_{X_k} + \Delta_k))$$

is generated in degree 1, so that there is a surjection

$$S^l H^0(m(K_{X_k} + \Delta_k)) \twoheadrightarrow H^0(ml(K_{X_k} + \Delta_k)).$$

Consider the following diagram:

$$\begin{array}{ccc}
 S^l H^0(m(K_X + \Delta)/R) & \xrightarrow{\alpha} & H^0(ml(K_X + \Delta)/R) \\
 \downarrow & & \downarrow \\
 S^l H^0(m(K_{X_k} + \Delta_k)) & \xrightarrow{\rightarrow} & H^0(ml(K_{X_k} + \Delta_k))
 \end{array}$$

The vertical maps are surjective by Theorem 17 (and applying Nakayama's lemma and cohomology and base change to pull back the generators of the k -modules). Now applying the version of Nakayama's Lemma given in [GH14, Chapter 5.3], surjectivity of the bottom map implies surjectivity of α , and thus $R(K_X + \Delta)$ is finitely generated over R . \square

Proposition 19. [Has15] *Let X be a two dimensional \mathbb{Q} -factorial normal variety over an algebraically closed field of characteristic $p > 0$. Let $\{(X, \Delta_i)\}_{i \in \{1, \dots, k\}}$ be a big, log smooth, \mathbb{Q} -Cartier KLT pair for each i and such that $K_X + \sum \Delta_i$ has simple normal crossings support. Then the ring*

$$R(X, K_X + \Delta_1, \dots, K_X + \Delta_k)$$

is finitely generated.

Corollary 20. *Let $\{(X, \Delta_i)\}_{i \in \{1, \dots, n\}}$ be a collection of klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta_i$ is big, \mathbb{Q} -Cartier and such that $K_X + \sum \Delta_i$ has simple normal crossings support. Then the ring*

$$R(X, K_X + \Delta_1, \dots, K_X + \Delta_n)$$

is finitely generated over R .

Proof. After base change, we may assume that R is complete and k is algebraically closed. C.f. the proof of [CZ12, 2.11], it suffices to prove that if $G = \sum_{i=1}^k m_i (a(K_X + \Delta_i))$, with fixed $a \in \mathbb{N}$, $m_1, \dots, m_k \geq 0$, and $l \in \{1, \dots, k\}$ then there exists m_l such that the following maps are surjective

$$\begin{aligned} H^0(X, \mathcal{O}_X(G - a(K_X + \Delta_l))) \otimes H^0(X, \mathcal{O}_X(a(K_X + \Delta_l))) \\ \rightarrow H^0(X, \mathcal{O}_X(G)). \end{aligned}$$

By applying Nakayama's Lemma and Theorem 17 as in the proof of Theorem 18, it suffices to show the surjectivity on the central fiber which holds by proposition 19. \square

8. FINITENESS OF MODELS FOR SURFACES

The following theorem is not anything new, but is stated here as the minimal model program is not usually run with scaling on a surface (in [KK94, T⁺14], termination is proven without scaling).

Theorem 21. *Let X be a two dimensional normal variety over a perfect field of characteristic $p > 0$. Let (X, Δ) be a log smooth KLT pair with a good minimal model (which exists for non-negative Kodaira dimensions by [T⁺14]) and let A be an ample \mathbb{Q} -divisor on X . Then there exists an $\epsilon > 0$ such that the minimal models (and the output of the minimal model program with scaling) for $(X, \Delta + tA)$ are all isomorphic for $t \in [0, \epsilon]$.*

Proof. (Essentially same as [Lai11, 26,27] and simplifies due to the fact that a birational isomorphism in codimension 1 on a surface gives an isomorphism). Let $\phi : (X, \Delta) \rightarrow (X_g, \Delta_g)$ be a good minimal model. Let $f : X_g \rightarrow Z = \text{Proj } R(K_{X_g} + \Delta_g)$ the contraction (which exists by since $K_{X_g} + \Delta_g$ is nef, hence semiample by [T⁺14]). Then ϕ contracts the divisorial part of $\mathbb{B}(K_X + \Delta)$. Pick $t_0 > 0$ such that $(X_g, \Delta_g + t_0\phi_*A) = (X_g, \Delta_g + t_0A_g)$ is klt with A ample on X (note A_g is big, not in general nef). For H ample on X_g , let $\psi : X_g \rightarrow X'$ be the minimal model program with scaling, which terminates (again by [T⁺14]), and gives a minimal model of $(X_g, \Delta_g + t_0\Delta_g)$ over Z . For any curve contracted by f , $(K_{X_g} + \Delta_g) \cdot C = 0$, hence $K_{X'} + \psi_*\Delta_g = K_{X'} + \Delta' \equiv_Z 0$. Thus curves contracted by ψ have trivial intersection with $K_{X_g} + \Delta_g$, and intersect negatively with A_g . Thus changing t does not affect which curves intersect negatively with $K_{X_g} + \Delta_g + tA_g$, and so X' is a minimal model of $(X_g, \Delta_g + tA_g)$ for all $t \in (0, t_0]$.

Now $\Delta' + t_0A'$ with $A' = \psi_*A_g$ (which is big) implies by the Theorem 10, that there exist only finitely many $K_{X'} + \Delta' + t_0A'$ negative extremal rays in $\overline{NE}(X')$. These are all necessarily just intersecting A' negatively, so decreasing t_0 , eventually we get to a point where a further decrease in t_0 doesn't change the number of negative extremal rays. Pick such a t_0 (now we are on X' , not over Z). Again shrinking t_0 , suppose that $\psi \circ \phi$ is discrepancy negative w'r't $(X, \Delta + tA)$ for all t in the half open interval $(0, t_0]$. Note that $\mathbb{B}(K_X + \Delta + t_0A) \subset \mathbb{B}(K_X + \Delta)$,

so ψ contracts only the curves not contracted by ϕ . Thus $\psi \circ \phi$ is discrepancy negative on the closed interval $[0, t_0]$, so X' is a minimal model of $(X, \Delta + tA)$ for all $t \in [0, t_0]$. Thus $\mathbb{B}(K_X + \Delta + tA)$ has the same divisorial components for all $t \in [0, t_0]$. Hence, any two minimal models for different $t \in [0, t_0]$ are birational and isomorphic in codimension one, and as the total dimension is two, they are thus isomorphic. \square

9. BIG ABUNDANCE / R

Definition 22. Let D_1, \dots, D_r be \mathbb{Q} -Cartier \mathbb{Q} -divisors on X . D_i are called **adjoint divisors** on X if they are of the form $D_i = K_X + \Delta_i$ for some **pair** (X, Δ_i) where X is normal and projective, $\Delta_i \geq 0$ and is a \mathbb{Q} -divisor. Let

$$R = R(X; D_1, \dots, D_r) = \bigoplus_{(n_1, \dots, n_r) \in \mathbb{N}^r} H^0(X, n_1 D_1 + \dots + n_r D_r).$$

and define the **support** of R to be

$$\text{Supp } R = \left(\sum_{i=1}^r \mathbb{R}_+ D_i \right) \cap \text{Div}_{\mathbb{R}}^{\text{eff}}(X).$$

Theorem 23. [CL13, 3.5] *Let X be a normal scheme of relative dimension 2, proper over a DVR R which has perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Let D_1, \dots, D_r be \mathbb{Q} -Cartier divisors on X . Assume that $R(X; D_1, \dots, D_r)$ is finitely generated. Let $D : \mathbb{R}^r \ni (\lambda_1, \dots, \lambda_r) \mapsto \sum \lambda_i D_i \in \text{Div}_{\mathbb{R}}(X)$ be the tautological map.*

(1) *The support of R is a rational polyhedral cone.*

(2) If $\text{Supp } R$ contains a big divisor and $D \in \sum \mathbb{R}_+ D_i$ is pseudoeffective, then $D \in \text{Supp } R$.

(3) There is a finite rational polyhedral subdivision $\text{Supp } R = \bigcup C_i$ such that σ_Γ is a linear function on C_i for every geometric valuation Γ of X . Furthermore, there is a coarsest subdivision with this property, in the sense that, if i and j are distinct, there is at least one geometric valuation Γ of X such that (the linear extensions of) $\sigma_\Gamma|_{C_i}$ and $\sigma_\Gamma|_{C_j}$ are different.

(4) There is a finite index subgroup $\mathbb{L} \subset \mathbb{Z}^r$ such that for all $n \in \mathbb{N}^r \cap \mathbb{L}$, if $D(n) \in \text{Supp } R$, then $\sigma_\Gamma(D(n)) = \text{mult}_\Gamma |D(n)|$.

Proof. Parts (1) and (2) follow easily from the cited theorem. For (3) and (4) note that [ELM⁺06, 4.7] is stated for an arbitrary Noetherian scheme, and this theorem is the only result used by the proof given in [CL13, 3.5]. \square

Theorem 24. ([CL13, 3.6]) *Let (X, Δ) be a klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is \mathbb{Q} -Cartier and that $R(K_X + \Delta, K_X + \Delta + A)$ is finitely generated. If $K_X + \Delta$ is pseudoeffective, then it is \mathbb{Q} -effective.*

Proof. As in the source paper. \square

Let Γ a geometric valuation on X . Define

$$\sigma_\Gamma = \inf \{ \text{mult}_\Gamma D' \mid D \sim_{\mathbb{R}} D' \geq 0 \} \in \mathbb{R}.$$

As a result of Theorem 20, there is the following abundance theorem:

Corollary 25. [CL13, 3.8] *Let (X, Δ) be a big klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $R(K_X + \Delta)$ and $R(K_X + \Delta, K_X + \Delta + A)$ are finitely generated for an ample \mathbb{Q} -divisor A on X . If $K_X + \Delta$ is nef, then it is semi-ample.*

Proof. (This simplifies from the cited theorem as I assume the pair $K_X + \Delta$ is big). For any $\epsilon > 0$, and any geometric valuation Γ on X , as $K_X + \Delta + \epsilon A$ is ample, a positive multiple is basepoint free. Thus,

$$\sigma_\Gamma(K_X + \Delta + \epsilon A) = 0$$

so that σ_Γ is identically zero by Theorem 23(4). Therefore, the centre of Γ is not in $\mathbb{B}(K_X + \Delta)$ for any such Γ . \square

Corollary 26. *Let X be a normal projective scheme of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K and let D_1, \dots, D_r be \mathbb{Q} -Cartier \mathbb{Q} -divisors on X . Assume that the ring $R = R(X; D_1, \dots, D_r)$ is finitely generated, and let $\text{Supp } R = \bigcup C_i$ be a finite rational polyhedral subdivision such that for every geometric valuation Γ of X , σ_Γ is linear on C_i , as in Theorem 23. Denote $\varphi : \sum \mathbb{R}_+ D_i \rightarrow N^1(X)_\mathbb{R}$ the projection, and assume there exists k such that $C_k \cap \varphi^{-1}(\text{Amp } D) \neq \emptyset$. Then $C_k \subset \text{Supp } R \cap \varphi^{-1}(\text{Nef } X)$. If the subdivision is coarsest, then $C_k = \text{Supp } R \cap \varphi^{-1}(\text{Nef } X)$*

Proof. As in the source, but applying Theorems 23 and 25. \square

10. CONE, RATIONALITY, AND CONTRACTION / R

Definition 27. [CL13, def 4.1] Let W a finite dimensional real vector space, $C \subset W$ a closed convex cone spanning W , and $v \in W$. The visible boundary of C from v is

$$V = \{w \in \partial C \mid [v, w] \cap C = \{w\}\}.$$

Corollary 28. (c.f. [CL13, 4.2]) (*Kawamata's Rationality, Cone and Contraction Theorem*) Let (X, Δ) be a pair of relative dimension 2, proper over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $v_0 = [K_X + \Delta]$ is \mathbb{Q} -Cartier and is obtained from the log smooth klt pair $\phi : (X', \Delta') \rightarrow (X, \Delta)$ after a finite number of steps of the minimal model program with scaling associated to a big divisor A with $K_X + \Delta + tA$ nef on X/R for some $t > 0$. Let V be the visible boundary of $\text{Nef}(X/R)$ from $v_0 \in N^1(X)_{\mathbb{R}}$ and $u_0 \in V \cap (v_0, [K_X + \Delta + tA])$. Then there exists a locally polyhedral neighborhood $U \subset V$ of u_0 such that $\forall u' \in U$, u' is semiample.

Proof. This is essentially a local version of the cited theorem. After base change, we may assume R is complete and that k is algebraically closed. Let $\mathcal{C} = \mathbb{R}_+ v_0 + \text{Nef}(X/R)$. Let u'_0 be a rational point of $N^1(X/R)$ sufficiently close to u_0 such that $u_0 \in B(u'_0, \epsilon)$ is a sup-norm ball with rational vertices $w_j = [K_X + \Delta + A_j]$, $j = 1, \dots, r$ which are klt and such that there exist big, log smooth, klt pairs $K_{X'} + \Delta' + A'_j$ with $\phi_*(K_{X'} + \Delta' + A'_j) = K_X + \Delta + A_j$ on X' , and such that ϵ is small enough that the w_j pull back to divisors having log smooth support on X' . Let \mathcal{B} denote the convex hull of $B(u'_0, \epsilon)$. For ϵ sufficiently small,

steps of the $K_{X'} + \Delta'$ minimal model program with scaling of A' are also steps of the $K_X + \Delta + A'_j$ minimal model program. Therefore there is an isomorphism of rings

$$\begin{aligned} & R(X, K_X + \Delta + A_1, \dots, K_X + \Delta + A_r) \\ & \approx R(X', K_{X'} + \Delta' + A'_1, \dots, K_{X'} + \Delta' + A'_r) \end{aligned}$$

so that by Theorem 20, these rings are finitely generated. Let $\varphi : \sum \mathbb{R}_+(K_X + \Delta + A_j) \rightarrow N^1(X)_{\mathbb{R}}$ the natural projection so that

$$\varphi\left(\sum \mathbb{R}_+(K_X + \Delta + A_j)\right) = \mathbb{R}_+\mathcal{B}$$

. Theorem 23 implies that $\text{Supp } R$ is a rational polyhedral cone which intersects the interior of the nef cone. Therefore, by Theorem 23(2), if U is the portion of $\partial \text{Nef}(X/R)$ contained in $B(u'_0, \epsilon)$, then $\varphi^{-1}(U) \subset \text{Supp}(R)$. Let $\text{Supp } R = \bigcup \mathcal{L}_k$ be the coarsest subdivision given by Theorem 23(3). For some k , $\mathcal{L}_k \cap \varphi^{-1}(\text{Amp } X) \neq \emptyset$, so by Theorem 26, $\mathcal{L}_k = \text{Supp } R \cap \varphi^{-1}(\text{Nef}(X))$, and thus U is locally polyhedral. If $u' \in U$, with divisor D then there exists an ample \mathbb{Q} -divisor A' such that $D \sim_{\mathbb{Q}} K_X + \Delta + A'$. By choice of ϵ , there is a log smooth divisor D' on X' with $\phi_* D' = D$, and so that $R(K_X + \Delta + A', K_X + \Delta + A' + A'')$ and $R(K_X + \Delta + A')$ are finitely generated for any ample A'' such that $D + A'' \in B(u'_0, \epsilon)$. Thus, Theorem 25 gives that u' is semiample. \square

Lemma 29. [CL13, Lemma 5.1] *Let X and Y be \mathbb{Q} -factorial projective schemes of relative dimension 2, smooth over a DVR R with algebraically closed residue field. Let $f : X \rightarrow Y$ be a birational contraction,*

and let $\tilde{f} : \mathbb{K}(Y) \approx \mathbb{K}(X)$ the induced isomorphism on function fields.

Then:

- (1) $f_* \operatorname{div}_X \varphi = \operatorname{div}_Y f(\varphi)$ for every $\varphi \in \mathbb{K}(X)$;
- (2) for every geometric valuation Γ on $\mathbb{K}(X)$, and for every $\varphi \in \mathbb{K}(X)$, we have $\operatorname{mult}_\Gamma(\operatorname{div}_X \varphi) = \operatorname{mult}_\Gamma(\operatorname{div}_Y \tilde{f}(\varphi))$;
- (3) if f is an isomorphism in codimension one, then $f_* : \operatorname{Div}_{\mathbb{R}}(X) \rightarrow \operatorname{Div}_{\mathbb{R}}(Y)$ is an isomorphism, and for every $D \in \operatorname{Div}_{\mathbb{R}}(X)$, there is an isomorphism $H^0(X, D) \approx H^0(Y, f_* D)$.

Proof. Follows easily from the cited Theorem. \square

Lemma 30. [CL13, 5.2] *Let X and Y be proper of relative dimension 2, over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K and $f : X \rightarrow Y$ a birational map which is an isomorphism in codimension one. Let $\mathcal{C} \subset \operatorname{Div}_{\mathbb{R}}^{\operatorname{eff}}(X)$ be a cone, and fix a geometric valuation Γ of X . Then the asymptotic order of vanishing σ_Γ is linear on \mathcal{C} if and only if it is linear on $f_* \mathcal{C} \subset \operatorname{Div}_{\mathbb{R}}^{\operatorname{eff}}(Y)$.*

Proof. As in the cited theorem, substituting Lemma 29 as necessary. \square

Lemma 31. [CL13, 6.2] *Let (X, Δ) be a log smooth klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Suppose (X, Δ) is not nef, and A is a big \mathbb{Q} -divisor such that $(X, \Delta + A)$ is klt, and $K_X + \Delta + A$ is nef. Let λ be the nef threshold. Then $\lambda \in \mathbb{Q}^+$, and there is $R \subset \overline{NE}(X)$ with $(K_X + \Delta + \lambda A) \cdot R = 0$ and $(K_X + \Delta) \cdot R < 0$.*

Proof. As in the cited theorem, but substituting Theorems 26 and 20 for their corresponding versions. \square

11. NEGATIVITY LEMMA

Lemma 32. [HK00, 1.7] *Let $f : X \rightarrow Y$ and $g : X \rightarrow Z$ be birational contractions where all spaces considered are schemes of relative dimension 2 over a DVR with perfect residue field. Suppose $f^*(A) + E = g^*(B) + F$ for A ample, B nef, E f -exceptional, and F g -exceptional all divisors extending to both fibers. Then $f \circ g^{-1} : Z \rightarrow Y$ is regular.*

Proof. It suffices to apply the negativity lemma (the usual proof holds c.f. [KM08, 3.39]) so that $E = F$ and thus $f^*(A) = g^*(B)$. The result then follows from the rigidity lemma which also holds under the same proof as [KM08, 1.6]. \square

Lemma 33. [CL13, 6.4] *Let (X, Δ) a projective klt pair, and $f : X \rightarrow Y$ a composite of $(K_X + \Delta)$ -divisorial contractions and $(K_X + \Delta)$ -flips. Then for every resolution*

$$\begin{array}{ccc} W & \xrightarrow{p} & X \\ q \downarrow & \nearrow f & \\ Y & & \end{array}$$

of f ,

$$p^*(K_X + \Delta) = q^*(K_Y + f_*\Delta) + E$$

with $E > 0$ a q -exceptional divisor. Therefore f cannot be an isomorphism. By the above formula,

$$H^0(X, K_X + \Delta) \approx H^0(Y, K_Y + f_*\Delta).$$

Proof. Follows easily from [KM08, 3.38]. \square

Lemma 34. [Har77, Exc II.4.2] *Let X be a reduced scheme, Y a separated scheme, and let f and g be two morphisms from X to Y . Assume that $f|_U = g|_U$ on a Zariski dense open subset $U \subset X$. Then $f = g$.*

12. FLIPS

Definition 35. [KM08, 3.33] Let X a normal scheme and D a \mathbb{Q} -divisor on X such that $K_X + D$ is \mathbb{Q} -Cartier. A $(K_X + D)$ -**flipping contraction** is a proper birational morphism $f : X \rightarrow Y$ to a normal scheme Y such that $Exc(f)$ has codimension at least two in X and $-(K_X + D)$ is f -ample. A normal scheme X^+ together with a proper birational morphism $f^+ : X^+ \rightarrow Y$ is called a $(K + D)$ -**flip** of f if

- (1) $K_{X^+} + D^+$ is \mathbb{Q} -Cartier, where D^+ is the birational transform of D on X^+
- (2) $K_{X^+} + D^+$ is f^+ -ample, and
- (3) $Exc(f^+)$ has codimension at least two in X^+ .

The induced rational map $\phi : X \rightarrow X^+$ is sometimes called a $(K + D)$ -flip by abuse of notation.

Theorem 36. *Let (X, Δ) be a projective \mathbb{Q} -factorial klt big pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Suppose that $f : X \rightarrow Y$ is a $K + \Delta$ flipping contraction. If $R(K_X + \Delta)$ is finitely generated, then the $(K + \Delta)$ -flip of f exists.*

Proof. Follows easily as in [KM08, 6.2], [CL13, 6.3]. □

Remark 37. By [EKW04, 1.2], the flip of f in the situation above is \mathbb{Q} -factorial.

13. RUNNING THE MINIMAL MODEL PROGRAM FOR GENERAL TYPE

Theorem 38. *Let (X, Δ) be a projective \mathbb{Q} -factorial klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Suppose $K_X + \Delta$ is big and log smooth over R . Then the minimal model program for (X, Δ) can be run, resulting in a terminating ssequence of flips and divisorial contractions.*

Proof. If necessary, extend R to be complete with algebraically closed residue field k . I begin similar to [CL13, 6.2]. Let A be a big divisor such that $K_X + \Delta + A$ is nef. Let α_1 be the smallest positive real number such that $K_X + \Delta + \alpha_1 A$ is nef. Denote by $\varphi : \text{Div}_{\mathbb{R}}(X) \rightarrow N^1(X)_{\mathbb{R}}$ the natural projection, and let $\|\cdot\|$ any norm on $N^1(X)_{\mathbb{R}}$. Pick finitely many big \mathbb{Q} -divisors H^1, \dots, H^r such that:

$$(1) \|\varphi(\Delta + \alpha_1 A) - \varphi(H^i)\| \ll 1 \text{ for all } i$$

(2) writing $\mathcal{C} = \sum_{i=1}^r \mathbb{R}_+ (K_X + H^i) \subset \text{Div}_{\mathbb{R}}(X)$, we have $K_X + \Delta + \alpha_1 A \in \text{int } \mathcal{C}$, and the dimension of the cone $\varphi(\mathcal{C}) \subset N^1(X)_{\mathbb{R}}$ is $\dim N^1(X)_{\mathbb{R}}$

(3) (X, H_i) is klt and log smooth for all i with $\text{supp}(K_X + \sum H_i)$ simple normal crossings.

In the first step, since H_i are log smooth, then by Theorem 20, the ring

$$R(X, K_X + \Delta, K_X + H^1, \dots, K_X + H^r)$$

is finitely generated. Thus applying Theorem 26, $\text{Supp } R = \bigcup C_j$ is rational polyhedral and $C^j = \text{Supp } R \cap \varphi^{-1}(\text{Nef}(X))$ is rational polyhedral, and there is a rational codimension 1 face of $\text{Nef}(X)$. Thus $\alpha_1 \in \mathbb{Q}_+$, and we can find an extremal ray $\mathfrak{R} \subset \overline{NE}(X)$ dual to this face which will satisfy $(K_X + \Delta + \alpha_1 A) \cdot \mathfrak{R} = 0$ and $(K_X + \Delta) \cdot \mathfrak{R} < 0$. By Theorem 28, this ray can be contracted under a birational contraction: $f' : X \rightarrow X'$.

If f' is a divisorial contraction, write $f_1 = f'$ and set $\Delta_2 = f_{1*}\Delta$, $A_2 = f_{1*}A$, so A_2 is again big and $K_{X_2} + \Delta_2 + \alpha_1 A_2$ is nef. By properties 1 and 3 above we have finite generation of $R(X_2, K_{X_2} + \Delta_2)$, $R(X_2, K_{X_2} + \Delta_2, K_{X_2} + \Delta_2 + A_2)$, and

$$R(X_2, K_{X_2} + \Delta_2, K_{X_2} + H_2^1, \dots, K_{X_2} + H_2^r)$$

$$\approx R(X, K_X + \Delta, K_X + H^1, \dots, K_X + H^r)$$

so the argument in the previous paragraph gives another extremal ray and the process can be repeated.

If, on the other hand, f' is a small contraction, then by Theorem 36, and applying Theorem 18, the flip of the contraction exists (c.f. definition 35). Write f_1 for the induced morphism to the flip and set $\Delta_2 = f_{1*}\Delta$, $A_2 = f_{1*}A$, so A_2 is again big and $K_{X_2} + \Delta_2 + \alpha_1 A_2$ is nef. By Lemma 29, we have finite generation of

$$R(X_2, K_{X_2} + \Delta_2)$$

and

$$R(X_2, K_{X_2} + \Delta_2, K_{X_2} + \Delta_2 + A_2).$$

Finally, we can let $H_2^r = f_{1*}H^r$, so that applying Lemma 29 to achieve finite generation of

$$R(X_2, K_{X_2} + \Delta_2, K_{X_2} + H_2^1, \dots, K_{X_2} + H_2^r),$$

we can similarly repeat the process.

Now suppose there is an infinite sequence of flips starting at step j . I will show leads to a contradiction. I proceed almost verbatim as in [CL13, 6.5]. By construction, C^j contains an open neighborhood of the nef divisor $K_{X_j} + \Delta_j + \alpha_j A_j$, so C^j contains ample divisors in its interior. Thus the cone $\varphi(\text{Supp } R_j) \subset N^1(X_j)$ also has dimension $\dim N^1(X_j)_{\mathbb{R}}$. Let $\text{Supp } R^j = \bigcup_k C_k^j$ the coarsest finite rational polyhedral subdivision from Theorem 23(3). For $i > j$, let $C_k^i \subset \text{Div}_{\mathbb{R}}(X_i)$ denote the proper transform of C_k^j and $C^i \subset \text{Div}_{\mathbb{R}}(X_i)$, the proper transform of C^j . By Lemma 30, for every geometric valuation Γ , the asymptotic order function σ_{Γ} is linear on each C_k^j .

By construction if $0 < \alpha \leq \alpha_1$, then $K_{X_j} + \Delta_j + \alpha A_j \in \text{Int } C^j$, so $K_{X_i} + \Delta_i + \alpha_i A_i \in \text{int } C^i$ for all $i > j$. Since $K_{X_i} + \Delta_i + \alpha_i A_i$ is nef, then by requirement (4) above and Lemma 24, $K_{X_i} + \Delta_i + \alpha_i A_i \in \text{Supp } R_i$. Thus, by Theorem 26, for each i there exists an index k such that the image of C_k^i in $N^1(X_i)_{\mathbb{R}}$ is a subset of $\text{Nef}(X_i)$. Therefore,

$$\varphi(C_k^1) \subset (f_{i-1} \circ \cdots \circ f_1)^* \text{Nef}(X_i).$$

Since there are finitely many cones C_k^j , there are two indices p and q such that $(f_{p-1} \circ \cdots \circ f_1)^* \text{Nef}(X_p)$ and $(f_{q-1} \circ \cdots \circ f_1)^* \text{Nef}(X_q)$ share a common interior point. Thus, by Lemma 32, the map $X_p \rightarrow X_q$ is a morphism. By Lemma 34, it is therefore an isomorphism. This contradicts Remark 33. \square

14. REDUCTIONS TO GENERAL CASE

Lemma 39. [HMX14, 2.8.3] *Let (X, Δ) be a log smooth pair which is a scheme of dimension at most 3, with the coefficients of Δ belonging to $(0, 1]$, and with X projective. If (X, Δ) has a weak log canonical model, then there is a sequence $\pi : Y \rightarrow X$ of smooth blow ups of the strata of Δ such that if we write $K_Y + \Gamma = \pi^*(K_X + \Delta) + E$, where $\Gamma \geq 0$ and $E \geq 0$ have no common components, $\pi_*\Gamma = \Delta$ and $\pi_*E = 0$ and if we write*

$$\Gamma' = \Gamma - \Gamma \wedge N_{\sigma}(Y, K_Y + \Gamma),$$

then $\mathbb{B}_-(Y, K_Y + \Gamma')$ contains no strata of Γ' . If Δ is a \mathbb{Q} -divisor, then Γ' is a \mathbb{Q} -divisor.

Proof. Follows easily from the cited theorem. \square

Lemma 40. [HMX14, 5.3] *Let X/R a projective scheme of relative dimension 2 over a DVR R with algebraically closed residue field and perfect fraction field. Let (X, Δ) be a log smooth dlt pair with X \mathbb{Q} -factorial and projective and Δ a \mathbb{Q} -divisor. If Φ is a \mathbb{Q} -divisor such that*

$$0 \leq \Delta - \Phi \leq N_\sigma(X, K_X + \Delta),$$

then steps of the $K_X + \Phi$ -minimal model program are steps of the $K_X + \Phi + t(\Delta - \Phi)$ minimal model program when t is sufficiently small. Furthermore, termination for (X, Φ) implies termination for (X, Δ) .

Proof. (Similar proof to the cited Theorem). Let $f : X \rightarrow Y$ the result of running a $(K_X + \Phi)$ minimal model program and let $Y \rightarrow Z$ be the ample model of $K_X + \Phi$. Let $\Delta_t = t\Delta + (1 - t)\Phi$ so that if $0 < t \ll 0$, then f is also a $(K_X + \Delta_t)$ -MMP. Let C generate a $K_X + \Delta_t$ -negative extremal ray and suppose that $(K_X + \Phi) \cdot C > 0$. Applying Theorem 10 on the special fiber,

$$-4 \leq (K_X + \Delta) \cdot C < 0.$$

Then, following the inequalities in [HMX14, 5.1], gives a contradiction if $0 < t < \frac{1}{5}$. Thus every step of the $K_X + \Delta_t$ with scaling of an ample divisor is $(K_X + \Phi)$ -trivial, so after finitely many steps there is a model $g : X \rightarrow W$ contracting the components of $N_\sigma(X, K_X + \Delta_t)$ (this holds with no changes from [HMX14, 2.7.1]). Thus, as $\text{supp}(\Delta - \Phi) \subset \text{supp} N_\sigma(X, K_X + \Delta) = \text{supp} N_\sigma(X, K_X + \Delta_t)$, then $g_*(K_Y + \Phi)$ is nef and / or semiample whenever $g_*(K_X + \Delta)$ is, and [HMX14, 2.7.2] (which holds using only the log smooth resolution) implies that g is

a minimal model of (X, Δ) , which is good when (X, Φ) has a good minimal model. \square

Lemma 41. [GL13, 2.3] *Let X/R a scheme which is projective of relative dimension 2 over either a DVR R with algebraically closed residue field and perfect fraction field or over a field. Let (X, Δ) be a log smooth klt pair and $\varphi : W \rightarrow X$ a log resolution of (X, Δ) . Choose Δ_W so that $K_W + \Delta_W = \varphi^*(K_X + \Delta) + E$ with Δ_W and E effective \mathbb{Q} -Weil divisors with no common component. Let $F = \sum_{F_i \text{ } \varphi\text{-exceptional prime divisor}} F_i$ and $\Delta_W^\epsilon = \Delta_W + \epsilon F$, then (W, Δ_W^ϵ) is an ϵ -log smooth model of (X, Δ) . Then for $0 < \epsilon \ll 1$, a (good) minimal model $\phi : (W, \Delta_W^\epsilon) \rightarrow (W_{\min}, \Delta_{W_{\min}}^\epsilon)$ is also a (good) minimal model for (X, Δ) .*

Proof. Follows easily from [BCDHMcK09, 3.6.10, 3.6.11]. \square

15. EXISTENCE OF MINIMAL MODELS (FOR GERMS) SPECIAL CASE

Lemma 42. [HMX10, 3.1] *Let (X, Δ) be a \mathbb{Q} -factorial klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that*

$$\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$$

contains no non-canonical centres of (X_k, Δ_k) . Let $f : X \rightarrow Y$ be a step of the $(K_X + \Delta)$ -MMP. If f is birational and V is a non-canonical centre of (X, Δ) , then V is not contained in the indeterminacy locus of f , V_0 is not contained in the indeterminacy locus of f_0 , and the induced maps $\phi : V \rightarrow W$ and $\phi_k : V_k \rightarrow W_k$ are birational, where

$W = f(V)$. Let $\Gamma = f_*\Delta$. Then $\mathbb{B}_-(Y_0, K_{Y_0} + \Gamma_0)$ contains no non-canonical centres of (Y_0, Γ_0) (so we may repeat the process), and if V is a non-klt centre or $V = X$, then $\phi : V \rightarrow W$ and $\phi_0 : V_0 \rightarrow W_0$ are birational contractions. If f is a Mori fibre space, then f_0 is not birational.

Proof. (Follows easily from the cited Theorem). Suppose f is birational. Let V be a non-canonical centre of (X, Δ) . Let $g : X \rightarrow Z$ be the contraction of the extremal ray associated to f (so that $f = g$ unless f is a flip). Let $Q = g(V)$, and let $\psi : V \rightarrow Q$, be the induced morphism. As every component of V_k is a non-canonical centre of (X_k, Δ_k) , then by hypothesis, components of V_k are not contained in $\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$, thus ψ_k is defined at each such component, hence is birational. By upper-semicontinuity of the fibers of ψ , ψ is birational, and thus $\phi : V \rightarrow Q \rightarrow W$, and ϕ_k are both birational.

Now suppose V is a non-klt centre or $V = X$, the above holds in the first case, as non-klt centres are non-canonical. Comparing the discrepancies of the differentials of adjunction for Δ_k and Γ_k as in the cited proof shows that ϕ_k, f_k , and thus ϕ are birational contractions. On the other hand, if f is a Mori fibre space, then, as the dimension of the fibers of $f : X \rightarrow Y$ are upper-semicontinuous, f_k is not birational. \square

Theorem 43. [HMX14, 3.2] *Let (X, Δ) be a \mathbb{Q} -factorial klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$ contains no non-canonical centres of (X_k, Δ_k) and is log smooth. Let*

$f : X \rightarrow Y$ be a step of the $(K_X + \Delta)$ -MMP. Then the minimal model program with scaling terminates using only contractions.

Proof. (Similar proof to the cited theorem). Let $f : X \rightarrow Y$ the $(K_X + \Delta)$ -MMP with scaling of an ample divisor A . Let $\Gamma = f_*\Delta$ and $B = f_*A$. By construction, $K_Y + tB + \Gamma$ is nef for some $t > 0$. By Lemma 42, $f : X \rightarrow Y$ is a birational contraction and $f_k : X_k \rightarrow Y_k$ is a birational contraction from $(X_k, tA_k + \Delta_k)$. If $K_X + \Delta$ is not pseudo-effective, then for $t > 0$, the result of f is a Mori fiber space, and by Lemma 42, Y_k is covered by $K_{Y_k} + tB_k + \Gamma_k$ -negative curves, contradicting bigness of $K_{X_k} + tA_k + \Delta_k$. Thus $K_X + \Delta$ is pseudo-effective, and given any $\epsilon > 0$, we may run the MMP until $t < \epsilon$. Now we conclude by Theorem 21. Letting ϵ be the constant given in that theorem, the minimal models for $(X_k, tA_k + \Delta_k)$ are all isomorphic $t \in [0, \epsilon]$. Thus, once $t < \epsilon$, any additional step in the minimal model program with scaling must be an isomorphism on the special fiber, and thus an isomorphism. Thus there exists a minimal model (Y, Γ) for (X, Δ) . \square

16. EXISTENCE OF MINIMAL MODELS (FOR GERMS) GENERAL CASE

Theorem 44. *Let (X, Δ) be a klt pair of relative dimension 2, proper over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is \mathbb{Q} -Cartier, pseudo-effective, and log smooth over R . Then the minimal model of (X, Δ) exists.*

Proof. I again start by repeating the reduction of [HMX14, 6.1]. If necessary, extend R to be complete with algebraically closed residue field k , so the strata of Δ have irreducible fibers over R . Let $f_0 : Y_0 \rightarrow X_0$ be the birational morphism of Lemma 39. Under the log smooth hypothesis, and since the strata of Δ have irreducible fibers, and f_0 blows up strata of Δ_0 , we may extend f_0 to a birational morphism $f : Y \rightarrow X/R$ which is a composition of smooth blow ups of strata of Δ . Write

$$K_Y + \Gamma = f^*(K_X + \Delta) + E$$

with $\Gamma \geq 0$ and $E \geq 0$, $f_*\Gamma = \Delta$, and $f_*E = 0$. Then (Y, Γ) is log smooth and the fibres of components of Γ are irreducible. By Lemma 41, (X, Δ) has a minimal model if (Y, Γ_ϵ) has a good minimal model, where $(Y, \Gamma_\epsilon) = (Y, \Gamma + \epsilon F)$ is the ϵ -log smooth model of (X, Δ) with F the sum of f -exceptional divisors and $0 < \epsilon \ll 1$.

Replace (X, Δ) by (Y, Γ_ϵ) and set $\Theta_k = \Delta_k - \Delta_k \wedge N_\sigma(X_k, K_{X_k} + \Delta_k)$ so that $\mathbb{B}_-(X_k, K_{X_k} + \Theta_k)$ contains no strata of Θ_k . Let $0 \leq \Theta \leq \Delta$ be the unique divisor such that $\Theta_k = \Theta|_{X_k}$. Let H relatively ample such that $(X, \Theta + H)$ is log smooth over R , and $K_X + \Theta + H$ is big. There is a commutative diagram:

$$\begin{array}{ccc} \pi_* \mathcal{O}_X(m(K_X + \Theta) + H) & \longrightarrow & \pi_* \mathcal{O}_X(m(K_X + \Delta) + H) \\ \downarrow & & \downarrow \\ H^0(X_k, \mathcal{O}_{X_k}(m(K_{X_k} + \Theta_k) + H_k)) & \longrightarrow & H^0(X_k, \mathcal{O}_{X_k}(m(K_{X_k} + \Delta_k) + H_k)) \end{array}$$

with surjective columns by Theorem 17 and [LE02, 5.3.20(b)], and with the bottom row an isomorphism. Applying Nakayama's lemma gives an isomorphism on the top row, so that $\Theta \geq \Delta - \Delta \wedge N_\sigma(X, K_X + \Delta)$. Again applying Theorem 17, gives $\Theta = \Delta - \Delta \wedge N_\sigma(X, K_X + \Delta)$. Thus $\Delta - \Theta \leq N_\sigma(X, K_X + \Delta)$, so by Lemma 40, it suffices to find a minimal model for (X, Θ) . Replacing (X, Δ) by (X, Θ) , it suffices to assume that $\mathbb{B}_-(X_k, K_{X_k} + \Delta_k)$ contains no strata of Δ_0 . Letting A be an ample divisor, we run the minimal model program with scaling of A . Now the assumptions of Theorem 43 apply, so we know that (X, Θ) has a minimal model. \square

Part 2. Globalization

(Under construction) In the following sections, I attempt to globalize the results above to work over Dedekind Domains.

17. KLT VANISHING

In this section (which is still being improved), I study the computability of the constant given in the vanishing theorem in [Tan12, 2.11]. The point is to have some maximum constant for the entire family when we globalize Theorem 38. First a theorem of Terakawa:

Theorem 45. [Ter99] *Let X be a smooth projective surface over an algebraically closed field of characteristic $p > 0$ and let D be a big and nef Cartier divisor on X . Assume that either*

- (1) $\kappa(X) \neq 2$ and X is not quasi-elliptic with $\kappa(X) = 1$; or
- (2) X is of general type with

$p \geq 3$ and $(D^2) > \text{vol}(X)$ or

$p = 2$ and $(D^2) > \max\{\text{vol}(X), \text{vol}(X) - 3\chi(\mathcal{O}_X) + 2\}$.

Then

$$H^i(X, \mathcal{O}_X(K_X + D)) = 0$$

for all $i > 0$.

Recall the following covering Lemma:

Lemma 46. *Let X be an n -dimensional smooth variety. Let D be a \mathbb{Q} -divisor such that the support of the fractional part $\{D\}$ is simple normal crossing. Moreover, suppose that, for the prime decomposition $\{D\} = \sum_{i \in I} \frac{b^{(i)}}{a^{(i)}} D^{(i)}$, no integers $a^{(i)}$ are divisible by p . Then there exists a finite surjective morphism $\gamma : Y \rightarrow X$ from a smooth variety Y with the following properties:*

(1) *The field extension $K(Y)/K(X)$ is a Galois extension.*

(2) *γ^*D is a \mathbb{Z} -divisor (it seems the degree of the extension is an m clearing the denominators of the $D^{(i)}$.)*

(3) *$\mathcal{O}_X(K_X + \lceil D \rceil) \approx (\gamma_* \mathcal{O}_Y(K_Y + \gamma^*D))^G$, where G is the Galois group of $K(Y)/K(X)$.*

(4) *If D' is a \mathbb{Q} -divisor such that $\{D'\} = \{D\}$, then γ^*D' is a \mathbb{Z} -divisor, and $\mathcal{O}_X(K_X + \lceil D' \rceil) \approx (\gamma_* \mathcal{O}_Y(K_Y + \gamma^*D'))^G$.*

Remark 47. Several aspects of the proof of the above Lemma will be used in the following.

Applying the above to Tanaka's proof seems to give the following:

Theorem 48. ([Tan12, 2.6] *Weak Effective Kawamata-Viehweg Vanishing Theorem*) *Let X be a smooth projective surface over an algebraically closed field of characteristic $p \geq 0$. Let N be a nef and big \mathbb{R} -cartier and B a nef and big \mathbb{Q} -divisor whose fractional part is simple normal crossing, whose fractional part has no denominators divisible by the characteristic. Then there exists an r , computable in terms of the intersection numbers of the components of B, N , and K_X , such that*

$$H^i(X, \mathcal{O}_X(K_X + \lceil B \rceil + rN + N')) = 0$$

for every $i > 0$, and every nef \mathbb{R} -Cartier \mathbb{R} -divisor N' such that $rN + N'$ is a \mathbb{Z} -divisor.

Proof. (A slightly different proof than the cited Theorem). The fractional part of $B + rN + N'$ is equal to the fraction part of B when $rN + N'$ is a \mathbb{Z} -divisor. Thus we apply Lemma 46 to obtain a degree m cover $\gamma : Y \rightarrow X$ (independent of r and N') where Y is a smooth surface. Then

$$\begin{aligned} & H^1(K_X + \lceil B \rceil + rN + N') \\ &= H^1(K_X + \lceil B + rN + N' \rceil) \\ &= H^1(\gamma_*(K_Y + \gamma^*(B + rN + N'))^G) \\ &= H^1(Y, K_Y + \gamma^*B + r\gamma^*N + \gamma^*N')^G. \end{aligned}$$

the last term vanishing by Terakawa's Theorem and some $r \gg 0$, as by the proof of [KMM87, 1-1-1], the cover can be made general type.

Now I claim it is possible to compute an r such that the last term above is zero using the intersections and degrees of components of

$B = \sum a_i \Gamma_i$. It will suffice to show that Y is general type and that $(B + rN)^2 > K_Y^2$, so that Theorem 45 can be applied.

As B is big and nef, there exists an effective divisor D such that $B - \frac{1}{j}D \equiv A$, for $j \gg 0$, is ample. Following the proof of [KMM87, 1-1-1], let $M = kA$ be very ample (we can compute k by [DCF15]) and such that m clears the denominators of the components of B and such that $mM - \Gamma_i$ is very ample for all i (again we can compute such an m by [DCF15]: in the case that $\Gamma_i^2 < 0$, then ensure k is large enough so that $kA \cdot \Gamma_i > -\Gamma_i^2$, take $\Gamma'_i = \Gamma_i + M$ and by [DCF15, Theorem 1.2] find (computable) m' such that $m'M - \Gamma'_i = m''M - \Gamma_i$ is very ample. Replace m by m'' , and letting $H_i \in |mM - \Gamma_i|$ for all i (thus $\frac{1}{m}(\Gamma_i + H_i) \sim M$ is very ample), we then have (again c.f. the proof of [KMM87, 1-1-1])

$$\begin{aligned} & K_Y \\ &= \tau^* K_X + (m-1) \left(\sum (\tau^* \Gamma_i)_{red} + \sum (\tau^* H_k^{(i)})_{red} \right) \\ &= \tau^* K_X + \tau^* \left(\sum_i \left(\Gamma_i + H_i - \frac{1}{m} (\Gamma_i + H_i) \right) \right) \\ &= \tau^* K_X + \tau^* ((m-1)M) \end{aligned}$$

so that Y is general type, and $vol(Y) = [((m+1)k+1)A]^2$, so it suffices to pick r large enough that $(B + rN)^2 > [((m+1)k+1)A]^2$ or, taking $j \gg 0$, such that $(B + rN)^2 > [((m+1)k+1)B]^2$ (or possibly a bit larger if $p = 2$ according to Theorem 45 hypothesis). \square

Now I restate Tanaka's vanishing theorem, with a note on the computability in certain circumstances.

Theorem 49. (*Effective KLT Kawamata-Viehweg Vanishing for normal surfaces c.f. [Tan12, 2.11]*) *Let (X, Δ) be a normal projective klt surface over an algebraically closed field of characteristic $p \geq 2$, where $K_X + \Delta$ is an effective \mathbb{R} -divisor. Let N be a nef and big \mathbb{R} -cartier \mathbb{R} -divisor. Let D be a \mathbb{Q} -Cartier \mathbb{Z} -divisor such that $D - (K_X - \Delta)$ is nef and big. Then there exists an r_0 ² such that*

$$H^i(X, \mathcal{O}_X(D + rN + N')) = 0$$

for every $i > 0$, every positive real number $r \geq r_0$, and every nef \mathbb{R} -Cartier \mathbb{R} -divisor N' such that $rN + N'$ is a Cartier divisor.

Proof. The computability follows easily from [Tan12, 2.11] except that Weak Kawamata Viehweg Vanishing in the proof is replaced by Theorem 48. Since, in this note I can apply this theorem when X is smooth and $\lfloor \Delta \rfloor = 0$, here is the proof in that simple case:

$$\begin{aligned} & H^1(D + rN + N') \\ &= H^1(K_X + D - K_X - \lfloor \Delta \rfloor + rN + N') \\ &= H^1(K_X + D - K_X + \lceil -\Delta \rceil + rN + N') \\ &= H^1(K_X + \lceil D \rceil - K_X - \Delta^\lceil + rN + N') \end{aligned}$$

²By Theorem 48 r is computable in terms of the number of components Δ and in terms of the self intersections of these components, and their multiplicities. Otherwise Tanaka's original statement gives just the existence of an r .

since we can move the integral divisor $D - K_X$ into the round up. This last term vanishes by Theorem 48. The vanishing of H^2 follows easily using Serre Duality. Otherwise if X is normal, one can keep track of the extra components of Δ gained in blowing up, and compute from that. \square

Now applying the same proof as in section 17 (where the same is proved but without a computable constant), we have the following:

Corollary 50. *Let (X, Δ) be a klt pair of relative dimension 2 over a DVR R with perfect residue field k of characteristic $p > 0$ and perfect fraction field K . Assume that $K_X + \Delta$ is big, \mathbb{Q} -Cartier, and simple normal crossings over R . Then, there exists a computable m_0 (depending only on the intersections components of K_X and Δ and possibly the picard number of X) such that for $m \gg 0, m \in m_0\mathbb{Z}^+$,*

$$h^0(m(K_{X_k} + \Delta_k)) = h^0(m(K_{X_K} + \Delta_K)).$$

18. GENERALIZATION OF TERAKAWA'S BASEPOINT FREE THEOREM

The first goal of this note is to give a very specific case of Theorem 2.4 from [Ter99] to KLT pairs:

Theorem 51. *Let (X, Δ) a minimal simple normal crossings projective KLT pair of general type and dimension 2 defined over an algebraically closed field of characteristic $p \geq 2$. Let $L = m(K_X + \Delta)$ be a nef and big line bundle on X . Assume that $l = L^2 \geq K_X^2 - 3\chi(\mathcal{O}_X) + 2 + 4d + 5$*

for $d \geq 0$. Then there is a computable constant m_0 (depending only on the intersections K_X and Δ and the number of components of Δ) such that $mm_0(K_X + \Delta)$ is basepoint-free.

The key lemma to the above is the following which follows easily from the cited paper.

Lemma 52. [Ter99, 2.2] *Let (X, Δ) a minimal simple normal crossings projective KLT pair of general type and dimension 2 defined over an algebraically closed field of characteristic $p \geq 2$. Let Z be a 0-cycle with $\deg Z = d+1$ where d is a nonnegative integer. Assume that L is a line bundle such that $K_X + \Delta + L$ is Cartier and $H^1(K_X + \Delta + L) = 0$. Assume that $K_X + L$ is $(d-1)$ -very ample and the restriction map is not surjective:*

$$\Gamma(K_X + \Delta + L) \rightarrow \Gamma(\mathcal{O}_Z(K_X + \Delta + L)).$$

Then there exists a rank 2 locally free sheaf E on X which is given by the short exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow I_Z(L) \rightarrow 0$$

where I_Z is the ideal sheaf of Z .

Proof of Theorem 51. Let m_0 be large enough so that the hypothesis of Theorem 49 holds i.e. such that

$$H^1(K_X + \Delta + L) = H^1(m_0(K_X + \Delta)) = 0.$$

If $D = K_X + \Delta + L$ is not base-point free, then there is a 0-cycle Z of degree 1 such that the restriction map

$$\Gamma(X, D) \rightarrow \Gamma(Z, D|_Z)$$

is not surjective. By Lemma 52, there exists a rank 2 locally free sheaf E on X which is given by the short exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow I_Z(L) \rightarrow 0$$

where I_Z is the ideal sheaf of Z . The rest of the argument follows exactly as in the cited Theorem.

19. MINIMAL MODELS FOR ARITHMETIC THREEFOLDS

Lemma 53. *Let (X, Δ) be a big log smooth klt pair of relative dimension 2 over a Dedekind Domain A . Assume that $K_X + \Delta$ is pseudo-effective and \mathbb{Q} -Cartier. Then for every $a \in A$*

$$N_\sigma(X, K_X + \Delta)|_{X_a} = N_\sigma(X_a, K_{X_a} + \Delta_a).$$

Proof. As in [HMX14, 2.3.1], it suffices to pick an ample A and show that there exists an m_0 such that for all m we have

$$f_*\mathcal{O}_X(mm_0(K_X + \Delta) + A) \twoheadrightarrow H^0(X_a, \mathcal{O}_{X_a}(mm_0(K_X + \Delta) + A))$$

for $m \gg 0$. As in [Kaw94, 5.8], we may assume that A is a complete discrete valuation ring whose residue field is algebraically closed. Applying [LE02, V.3.20], and the asymptotic deformation invariance of

plurigenera proved in section 17, it holds that

$$\begin{aligned} & H^0(X_a, \mathcal{O}_{X_a}(mm_0(K_X + \Delta) + A)) \\ & \approx H^0(X, \mathcal{O}_X(mm_0(K_X + \Delta) + A)) \otimes k(a) \end{aligned}$$

with $k(a) \approx A/\mathfrak{m}$, and so the lemma follows. \square

Now assume that (X, Δ) is general type. By the above Lemma, after blowing up to a log-smooth terminal family (as in 17) the divisor Θ constructed there has the same data along the fibers (self intersection and intersection with K_{X_a}) since by log-smoothness

$$\begin{aligned} \Theta_a &= \Delta_a - \Delta_a \wedge N_\sigma(K_{X_a} + \Delta_a) \\ &= \Delta|_{X_a} - \Delta|_{X_a} \wedge N_\sigma(K_X + \Delta)|_{X_a} \\ &= [\Delta - \Delta \wedge N_\sigma(K_X + \Delta)]|_{X_a}. \end{aligned}$$

After localizing at any point $a \in A$, the situation of Theorem 38 applies. In particular we have on $A_{(a)}$ all the necessary finite generations and termination of the minimal model program. As the picard number is bounded over A , and since the constant given by Tanaka's vanishing Theorem 6 only depends on the nef and big divisors used, we can find an r such that for all such a vanishing of

$$H^1\left(m(K_{X_{(a)}^g} + \Delta)\right)$$

holds, where $X_{(a)}^g$ is the good minimal model of X localized at a . After a base-change from $A_{(a)}$ to a complete DVR with algebraically closed residue field, the computable base-point free theorem 51. Applying the

proof of [CZ12, 2.7], there is thus a degree bound over all of A for the finite generations of the following rings:

$$R(K_X + \Delta/A)$$

$$R(K_X + \Delta_1, K_X + \Delta_2, \dots, K_X + \Delta_n/A)$$

where Δ_i and Δ are such that $K_X + \Delta, K_X + \Delta_i$ are big log smooth klt and $K_X + \sum \Delta_i$ has log-smooth support and Δ_i are sufficiently close to Δ for some norm on $N^1(X/A)$. Thus the following holds using the same proof (but substituting the above finite generations) as in Theorem 44:

Theorem 54. *Let (X, Δ) be a projective \mathbb{Q} -factorial klt pair of relative dimension 2 over a dedekind domain A . Suppose $K_X + \Delta$ is big and log smooth over R . Then the minimal model program for (X, Δ) can be run, resulting in a terminating sequence of flips and divisorial contractions.*

REFERENCES

- [Art69] Michael Artin. Algebraization of formal moduli. i. *Global analysis (papers in honor of K. Kodaira)*, pages 21–71, 1969.
- [Art70] Michael Artin. Algebraization of formal moduli: II. existence of modifications. *Annals of Mathematics*, pages 88–135, 1970.
- [BCDHMcK09] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. *Journal of the American Mathematical Society*, 23(2):405, 2009.

- [BCZ15] Caucher Birkar, Yifei Chen, and Lei Zhang. Iitaka’s $c_{\{n,m\}}$ conjecture for 3-folds over finite fields. *arXiv preprint arXiv:1507.08760*, 2015.
- [Bir15] Caucher Birkar. Existence of flips and minimal models for 3-folds in char p . *arXiv preprint arXiv:1311.3098*, 2015.
- [BLR12] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud. *Néron models*, volume 21. Springer Science & Business Media, 2012.
- [CL13] Alessio Corti and Vladimir Lazić. New outlook on the minimal model program, ii. *Mathematische Annalen*, 356(2):617–633, 2013.
- [CP14] Vincent Cossart and Olivier Piltant. Resolution of singularities of arithmetical threefolds ii. *arXiv preprint arXiv:1412.0868*, 2014.
- [CZ12] Paolo Cascini and DEQI ZHANG. Effective finite generation for adjoint rings. *arXiv preprint arXiv:1203.5204*, 2012.
- [DCF15] Gabriele Di Cerbo and Andrea Fanelli. Effective matsusaka’s theorem for surfaces in characteristic p . *arXiv preprint arXiv:1501.07299*, 2015.
- [Eke88] Torsten Ekedahl. Canonical models of surfaces of general type in positive characteristic. *Publications Mathématiques de l’IHÉS*, 67:97–144, 1988.
- [EKW04] E Javier Elizondo, Kazuhiko Kurano, and Kei-ichi Watanabe. The total coordinate ring of a normal projective variety. *Journal of Algebra*, 276(2):625–637, 2004.
- [ELM⁺06] Lawrence Ein, Robert Lazarsfeld, Mircea Mustață, Michael Nakamaye, and Mihnea Popa. Asymptotic invariants of base loci. In *Annales de l’institut Fourier*, volume 56, pages 1701–1734, 2006.
- [GH14] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. John Wiley & Sons, 2014.

- [GL13] Yoshinori Gongyo and Brian Lehmann. Reduction maps and minimal model theory. *Compositio Mathematica*, 149(02):295–308, 2013.
- [Gro67] A Grothendieck. *Éléments de géométrie algébrique*. New York, 1967.
- [Har77] Robin Hartshorne. *Algebraic geometry*, volume 52. Springer Science & Business Media, 1977.
- [Has15] Kenta Hashizume. Finite generation of adjoint ring for log surfaces. *arXiv preprint arXiv:1505.02282*, 2015.
- [HK00] Yi Hu and Sean Keel. Mori dream spaces and git. *arXiv preprint math/0004017*, 2000.
- [HMX10] Christopher Hacon, James McKernan, and Chenyang Xu. On the birational automorphisms of varieties of general type. *arXiv preprint arXiv:1011.1464*, 2010.
- [HMX14] CHRISTOPHER D Hacon, James McKernan, and Chenyang Xu. Boundedness of moduli of varieties of general type. *arXiv preprint arXiv:1412.1186*, 2014.
- [HX13] Christopher D Hacon and Chenyang Xu. On the three dimensional minimal model program in positive characteristic. *arXiv preprint arXiv:1302.0298*, 2013.
- [Kaw94] Yujiro Kawamata. Semistable minimal models of threefolds in positive or mixed characteristic. In *J. Algebraic Geom.* Citeseer, 1994.
- [Kaw98] Yujiro Kawamata. Index 1 covers of log terminal surface singularities. *arXiv preprint math/9802044*, 1998.
- [KK94] János Kollár and Sándor Kovács. Birational geometry of log surfaces. *preprint*, 1994.
- [KM08] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134. Cambridge University Press, 2008.

- [KMM87] Yujiro Kawamata, Katsumi Matsuda, and Kenji Matsuki. Introduction to the minimal model problem. *Adv. Stud. Pure Math*, 10:283–360, 1987.
- [Kol13] János Kollár. *Singularities of the minimal model program*, volume 200. Cambridge University Press, 2013.
- [KU85] Toshiyuki Katsura and Kenji Ueno. On elliptic surfaces in characteristic p . *Mathematische Annalen*, 272(3):291–330, 1985.
- [Lai11] Ching-Jui Lai. Varieties fibered by good minimal models. *Mathematische Annalen*, 350(3):533–547, 2011.
- [Laz04] Robert K Lazarsfeld. *Positivity in algebraic geometry I: Classical setting: line bundles and linear series*, volume 48. Springer Science & Business Media, 2004.
- [LE02] Qing Liu and Reinie Erne. *Algebraic geometry and arithmetic curves*. Oxford university press, 2002.
- [MP05] Yuri Ivanovic Manin and Alexei A Panchishkin. *Introduction to modern number theory*, volume 49. Springer, 2005.
- [Suh08] Junecue Suh. Plurigenera of general type surfaces in mixed characteristic. *Compositio Mathematica*, 144(05):1214–1226, 2008.
- [T⁺14] Hiromu Tanaka et al. Minimal models and abundance for positive characteristic log surfaces. *Nagoya Mathematical Journal*, 216:1–70, 2014.
- [Tan12] Hiromu Tanaka. The x-method for klt surfaces in positive characteristic. *arXiv preprint arXiv:1202.2497*, 2012.
- [Ter99] Hiroyuki Terakawa. The d-very ampleness on a projective surface in positive characteristic. *Pacific J. of Math*, 187:187–198, 1999.